

Boerboel Genetic Evaluation

SA Stud Book
February 2026

MR M. MOTLHAGODI

PO BOX 293

0

TSABONG

BOTSWANA



Kennel Number: 693999



Estimated Breeding Values (EBVs)

Litter size and conformation traits are influenced by many genes, with every gene making a small contribution to the expression of the trait. These traits are, however, also influenced by physiological and physical environmental factors, for example the age and sex of the animal, the feeding and management the animal received and year x season effects. It is therefore not possible to know what the animal's genetic potential for the trait is by merely using the measurement.

With BLUP (*Best Linear Unbiased Prediction*) methodology an animal's genetic potential is predicted by making use of the genetic parameters (heritabilities and genetic correlations) of the breed, pedigree information, as well as the measurements of the animal as deviation from the average of the animals that were measured with the animal. Adjustments are made for differences in sex, age, classifier, management and year x season effects to ensure that fair comparisons are made amongst the animals. These predictions are called **EBVs (Estimated Breeding Values)** which indicates the animal's genetic potential for the specific trait and predicts the type of progeny that the animal will breed.

Pedigree information determines the genetic level at which the animal enters the genetic evaluation, it links the information of all relatives together and it is used to determine the genetic ranking of the breed. For young animals that were not yet measured, mid-parent EBVs are estimated, which are the average of the EBVs of the sire and dam. These breeding values are still inaccurate and can change a lot when more information (measurements) is included in subsequent genetic evaluations.

Breeding values therefore enables a breeder to compare his/her animals genetically to one another, as well as to all other animals of the breed. This Report gives a summary of the genetic levels of the active males and females of the relevant breeder in comparison to that of the breed for linearly classified traits and litter size. The genetic change that occurred over time can also easily be seen.

Breeding Value Indices (EBVi)

For easy interpretation, EBVs are standardised and expressed relative to the average (100) of the active animals of the breed, using 12 as standard deviation (SD). This means that an EBV of 100 is equal to the active average of the breed, and more or less 50% of the animals will have EBVi's of between 90 and 110, which indicates an average animal. Only 0.5% of the EBVi's will be lower than 64 and 0.5% will be higher than 136. Higher EBVi's indicate better genetic potential for the trait.

Accuracy

EBVs are published with an **accuracy value (%)**, which indicates how much info was available for the estimation of the EBV and with how much trust it can therefore be applied. Factors that influence the accuracy include heritability, completeness and correctness of pedigree information, number of own and measured relatives and the size of the group in which the animal was measured. When an accuracy of 85% and more is reached, the EBV should not change significantly with the inclusion of more information in subsequent genetic evaluations.

Kennel and Breed Averages: The different traits' averages of the breeder's active animals are compared to those of the breed in the Kennel and Breed Average Graphs and Tables. The breeder can therefore easily see for which traits the animals are above or below the average of the breed.


Genetic Trends: The Genetic Trends indicate the genetic change that occurred over time in the animals bred by the breeder compared to that of the breed. These are calculated as the average of the breeding values of all measured animals bred by the breeder, per year of birth.

Inbreeding: Inbreeding is the mating of related animals and is widely viewed as being harmful. The danger of inbreeding lies there in that it can gradually decrease fertility and survivability – a phenomenon known as inbreeding depression. It can cause lower conception rates, more abortions, more stillborn and weak offspring and a higher susceptibility to diseases. Indications in dogs include a low litter size as well as problems relative to fertility, such as invisible heat, behavioural problems when mating and low sperm quality. Overall, reduced litter size most likely reflects inbreeding depression in several traits, both physiological and behavioural, as well as a high frequency of non-viable embryos due to lethal alleles. Generally, the effects of inbreeding can become noticeable at a value of 6.25%, therefore it is generally recommended that individuals should have inbreeding coefficients of less than 6.25%. Breeders should therefore avoid matings causing inbreeding of more than 6.25% in the progeny. The theoretical maximum inbreeding coefficient is 50%. However, it should be noted that not all inbred animals show signs of inbreeding depression.

The data used in this evaluation was obtained from LOGIX from linear classifications that were done by the Breeder Society. All possible attempts have been made to present the information correctly. SA Stud Book does not accept responsibility for the use and interpretation of the information presented on animals in this publication.

Genetic Levels

Kennel Average relative to Breed Average, where Breed Average is indicated as 0



Average Breeding Value Indices (EBVi) where Breed Average = 100

Trait	All Animals	Males	Females
Litter size (LS)	91	94	82
Movement (Mv)	91	91	89
Temperament (Tmp)	100	97	110
Coat (Ct)	98	98	96
General Appearance	103	98	123
Head & Face (H&F)	97	92	116
Front Quarter (FQ)	102	100	113
Centre (C)	109	107	119
Hind Quarter (HQ)	103	103	105
Final Score (FS)	101	98	113
Hip Health (HH)	86	89	77
Elbow Health (ELM)	104	105	101

Inbreeding

Notes

Inbreeding primarily leads to decreased fertility, low survivability of pups and low disease resistance.

Inbreeding coefficients are calculated for all animals of the breed and indicates the probability that two genes of a pair are the same due to a common ancestor in the pedigree of the animal.

The adverse effects of inbreeding can be observed from 6.25% onwards.


It is therefore highly recommended to record pedigree information as accurately. Inbreeding is dependent on pedigree information. Conventional methods for the calculation of inbreeding coefficients usually assume that unknown parents are unrelated, which is usually far from the truth. Inbreeding coefficients are therefore conservative indicators of the level of inbreeding.

Incomplete or incorrect pedigree information can be reported to SA Stud Book via the Breed Society.

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Rate of Inbreeding




Inbreeding Levels


	All Animals		Males		Females		Sires		Dams	
	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed
Number	5	49007	4	25141	1	23857	5	3371	5	6287
Average Inbreeding	10.08	8.49	10.85	8.46	6.99	8.53	14.75	8.01	12.70	8.00
Number inbred	5	48727	4	24990	1	23728	5	24990	5	23728
Number >6.25% Inbr.	5	30600	4	15537	1	15058	5	15537	5	15058

Genetic Trends


Litter Size




Movement




Temperament




Coat



General Appearance




Head & Face




Genetic Trends


Front Quarter



Centre




Hind Quarter




Final Score & Health Traits


Final Score



Hip Health



Elbow Health



Estimated Breeding Value Indices (EBVi's)

Males

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
JOSMOT TOP		TARGUS BOM BOEL	7.35	94	106	90	86	107	93	93	96	107	113	109	103	4	1	3			1
	2020	TREBLA MINNIE		5	5	33	57	34	70	56	69	58	60	62	66						1
BLCKOPS BAZOOKA		BLCKOPS POTTER	9.63	97	94	92	78	96	92	107	97	104	109	103	99	11	4	3			
	2021	JOSMOT LARA CROFT		5	5	5	37	5	57	36	56	38	40	44	50						
JOSMOT JAMES		ELDORADO BBD FRANK SINATRA	17.62	80	115	101	106	93	109	91	79	93	97	101	91						
	2023	JOSMOT KOOLKAT 2		5	5	17	49	18	65	48	64	50	53	55	60						
MOET KENNEL GUN		BLCKOPS BAZOOKA	8.79	84	105	93	94	93	99	100	97	94	109	100	99						
	2024	JOSMOT HANNAH		5	5	5	34	5	55	33	54	35	38	41	48						

Estimated Breeding Value Indices (EBVi's)

Females

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
JOSMOT THE EYE		MES HITLER BLAKE	6.99	77	101	82	89	110	96	123	116	113	119	105	113	4	2				
	2021	JOSMOT MEGA		5	5	31	47	19	63	46	62	48	50	52	58						

Estimated Breeding Value Indices (EBVi's)

Sires

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Sire Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels	Hips	Elbows		
TARGUS BOM BOEL		CABARET SON OF A GUN	24.74	81	114	66	90	114	79	112	94	103	116	107	102	86 14	38 28	3 3	28 8		
	2016	CRIHATA LILLIE		5	5	57	78	57	88	78	87	79	80	82	85						
ELDORADO BBD FRANK SINATRA		CABARET PAVAROTTI	18.64	65	125	98	113	88	107	96	83	89	96	99	91	51 10	28 20		12 4		
	2017	CABARET PAMELA ANDERSON		5	5	41	69	43	83	68	82	70	72	74	79						
MES HITLER BLAKE		MES MASCOT	13.12	78	100	91	101	121	98	125	119	126	134	114	123	37 6	18 17	1 1	8 1		
	2018	MES SERABI		27	46	42	66	44	79	65	78	67	69	71	75						
BLCKOPS POTTER		JOSMOT TATANKA	7.60	102	93	96	88	86	86	86	76	84	89	103	84	4 1	1 1		1 1		
	2019	ERNYMOS LATANSHA		5	5	5	35	5	50	35	49	36	38	41	46						
BLCKOPS BAZOOKA		BLCKOPS POTTER	9.63	97	94	92	78	96	92	107	97	104	109	103	99	11 4	3 2				
	2021	JOSMOT LARA CROFT		5	5	5	37	5	57	36	56	38	40	44	50						

Estimated Breeding Value Indices (EBVi's)

Dams

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe-r ament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
TREBLA MINNIE	2018	MAWEZA DISSELBOOM	10.18	107	98	119	89	101	110	89	108	114	110	105	108	20	4	12	1	1	10
		TREBLA ULA II		5	5	43	63	37	77	62	77	64	66	69	73			9			
JOSMOT MEGA	2019	CABARET BORIS (PASELLA)	11.73	76	102	83	75	95	81	112	99	94	101	93	94	11	2	2			1
		TARGUS NOVA		5	5	39	54	28	69	53	68	54	57	59	64			2			
JOSMOT LARA CROFT	2020	MES HITLER BLAKE	10.79	92	95	88	83	108	93	118	116	120	123	102	113	4	1	1			1
		TREBLA JASMYN		5	5	20	43	16	55	42	54	43	45	47	51			1			
JOSMOT KOOLKAT 2	2020	TARGUS BOM BOEL	7.35	94	106	103	90	104	102	92	92	109	107	107	101	23	2	7	1	1	3
		TREBLA MINNIE		5	5	44	61	36	75	59	74	62	64	66	70			5			
JOSMOT HANNAH	2021	TARGUS MOSCOW	23.45	71	116	94	100	98	94	100	92	91	109	99	97	5	1	1			
		TARGUS LUNA		5	5	24	38	5	51	38	51	39	41	43	47			1			

