

Boerboel Genetic Evaluation

SA Stud Book

February 2026

MR A. ROL

HAUWERT 3

0

1691 FA HAUWERT

NETHERLAND



Kennel Number: 706979



Estimated Breeding Values (EBVs)

Litter size and conformation traits are influenced by many genes, with every gene making a small contribution to the expression of the trait. These traits are, however, also influenced by physiological and physical environmental factors, for example the age and sex of the animal, the feeding and management the animal received and year x season effects. It is therefore not possible to know what the animal's genetic potential for the trait is by merely using the measurement.

With BLUP (*Best Linear Unbiased Prediction*) methodology an animal's genetic potential is predicted by making use of the genetic parameters (heritabilities and genetic correlations) of the breed, pedigree information, as well as the measurements of the animal as deviation from the average of the animals that were measured with the animal. Adjustments are made for differences in sex, age, classifier, management and year x season effects to ensure that fair comparisons are made amongst the animals. These predictions are called **EBVs (Estimated Breeding Values)** which indicates the animal's genetic potential for the specific trait and predicts the type of progeny that the animal will breed.

Pedigree information determines the genetic level at which the animal enters the genetic evaluation, it links the information of all relatives together and it is used to determine the genetic ranking of the breed. For young animals that were not yet measured, mid-parent EBVs are estimated, which are the average of the EBVs of the sire and dam. These breeding values are still inaccurate and can change a lot when more information (measurements) is included in subsequent genetic evaluations.

Breeding values therefore enables a breeder to compare his/her animals genetically to one another, as well as to all other animals of the breed. This Report gives a summary of the genetic levels of the active males and females of the relevant breeder in comparison to that of the breed for linearly classified traits and litter size. The genetic change that occurred over time can also easily be seen.

Breeding Value Indices (EBVi)

For easy interpretation, EBVs are standardised and expressed relative to the average (100) of the active animals of the breed, using 12 as standard deviation (SD). This means that an EBV of 100 is equal to the active average of the breed, and more or less 50% of the animals will have EBVi's of between 90 and 110, which indicates an average animal. Only 0.5% of the EBVi's will be lower than 64 and 0.5% will be higher than 136. Higher EBVi's indicate better genetic potential for the trait.

Accuracy

EBVs are published with an **accuracy value (%)**, which indicates how much info was available for the estimation of the EBV and with how much trust it can therefore be applied. Factors that influence the accuracy include heritability, completeness and correctness of pedigree information, number of own and measured relatives and the size of the group in which the animal was measured. When an accuracy of 85% and more is reached, the EBV should not change significantly with the inclusion of more information in subsequent genetic evaluations.

Kennel and Breed Averages: The different traits' averages of the breeder's active animals are compared to those of the breed in the Kennel and Breed Average Graphs and Tables. The breeder can therefore easily see for which traits the animals are above or below the average of the breed.


Genetic Trends: The Genetic Trends indicate the genetic change that occurred over time in the animals bred by the breeder compared to that of the breed. These are calculated as the average of the breeding values of all measured animals bred by the breeder, per year of birth.

Inbreeding: Inbreeding is the mating of related animals and is widely viewed as being harmful. The danger of inbreeding lies there in that it can gradually decrease fertility and survivability – a phenomenon known as inbreeding depression. It can cause lower conception rates, more abortions, more stillborn and weak offspring and a higher susceptibility to diseases. Indications in dogs include a low litter size as well as problems relative to fertility, such as invisible heat, behavioural problems when mating and low sperm quality. Overall, reduced litter size most likely reflects inbreeding depression in several traits, both physiological and behavioural, as well as a high frequency of non-viable embryos due to lethal alleles. Generally, the effects of inbreeding can become noticeable at a value of 6.25%, therefore it is generally recommended that individuals should have inbreeding coefficients of less than 6.25%. Breeders should therefore avoid matings causing inbreeding of more than 6.25% in the progeny. The theoretical maximum inbreeding coefficient is 50%. However, it should be noted that not all inbred animals show signs of inbreeding depression.

The data used in this evaluation was obtained from LOGIX from linear classifications that were done by the Breeder Society. All possible attempts have been made to present the information correctly. SA Stud Book does not accept responsibility for the use and interpretation of the information presented on animals in this publication.

Genetic Levels

Kennel Average relative to Breed Average, where Breed Average is indicated as 0



Average Breeding Value Indices (EBVi) where Breed Average = 100

Trait	All Animals	Males	Females
Litter size (LS)	91	92	90
Movement (Mv)	94	93	95
Temperament (Tmp)	92	95	90
Coat (Ct)	105	108	104
General Appearance	94	95	93
Head & Face (H&F)	99	104	96
Front Quarter (FQ)	98	103	95
Centre (C)	93	96	92
Hind Quarter (HQ)	96	99	94
Final Score (FS)	96	100	93
Hip Health (HH)	119	126	115
Elbow Health (ELM)	107	111	106

Inbreeding

Notes

Inbreeding primarily leads to decreased fertility, low survivability of pups and low disease resistance.

Inbreeding coefficients are calculated for all animals of the breed and indicates the probability that two genes of a pair are the same due to a common ancestor in the pedigree of the animal.

The adverse effects of inbreeding can be observed from 6.25% onwards.


It is therefore highly recommended to record pedigree information as accurately. Inbreeding is dependent on pedigree information. Conventional methods for the calculation of inbreeding coefficients usually assume that unknown parents are unrelated, which is usually far from the truth. Inbreeding coefficients are therefore conservative indicators of the level of inbreeding.

Incomplete or incorrect pedigree information can be reported to SA Stud Book via the Breed Society.

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Rate of Inbreeding




Inbreeding Levels


	All Animals		Males		Females		Sires		Dams	
	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed
Number	6	49007	2	25141	4	23857	5	3371	5	6287
Average Inbreeding	5.73	8.49	5.91	8.46	5.64	8.53	4.51	8.01	12.35	8.00
Number inbred	6	48727	2	24990	4	23728	5	24990	5	23728
Number >6.25% Inbr.	3	30600	1	15537	2	15058	1	15537	4	15058

Genetic Trends


Litter Size




Movement




Temperament




Coat



General Appearance




Head & Face




Genetic Trends


Front Quarter



Centre




Hind Quarter




Final Score & Health Traits


Final Score



Hip Health



Elbow Health



Estimated Breeding Value Indices (EBVi's)

Males

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe- rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels		Hips	Elbows	
IMMER MOED LAU		GIANTS OF AFRICA MATTEO	5.41	124	100	94	92	92	112	90	96	96	92	99	96	4	1		4	4	1
	2019	IMMER MOED USHI		5	23	5	14	5	29	13	28	15	17	20	25						
ALEWIJNS IMMERMUED NEO		AFRIKA LEONARDO III	6.41	128	121	91	95	97	103	99	111	110	100	99	105						
	2023	ALEWIJNS IMMERMUED BOMBARIE		5	20	5	29	5	46	28	45	30	32	35	40						

Estimated Breeding Value Indices (EBVi's)

Females

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
ALEWIJN'S IMMERMOED AMMEHOELA	2019	CHARMEUR FROM SOUS AFRICA (POETIN)	8.05	116	98	90	95	91	104	99	87	94	94	88	89	1		1			
		IMMERMOED NOSTRA DAME		28	51	5	5	5	15	5	13	5	5	5	5						
ALEWIJNS IMMERMOED BOMBARIE	2020	IMMER MOED LAU	3.85	115	102	87	90	87	108	86	99	93	86	95	93	4	1	1			
		BREKER HURRICANE HATTIE		5	13	5	17	5	30	16	30	17	20	22	26						
ALEWIJNS IMMERMOED ELA	2022	THE BONAFIDE GRONK	4.24	102	102	94	99	88	104	91	89	88	86	92	88						
		ALEWIJN'S IMMERMOED AMMEHOELA		5	14	5	20	5	39	19	38	21	24	27	33						
ALEWIJNS IMMERMOED TRAMMELANT	2023	AFRIKA LEONARDO III	6.41	128	121	91	95	96	102	97	109	107	99	101	104						
		ALEWIJNS IMMERMOED BOMBARIE		5	20	5	18	5	31	17	30	19	21	23	27						

Estimated Breeding Value Indices (EBVi's)

Sires

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Sire Information		Inbreed	Health		Fertility	General			Appearance						Progeny :	Appraisal	Health		Progeny with litters		
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters		Progeny Kennels	Hips	Elbows	
GIANTS OF AFRICA MATTEO		MIDDELPOS BBD ALPHA	3.76	121	105	93	103	103	104	102	106	113	110	119	112	2			2	2	1
	2015	GIANTS OF AFRICA MADONNA		26	33	47	51	39	57	51	57	52	52	53	55						1
THE BONAFIDE GRONK		KING OF THE CAGES DEUCE	3.83	88	106	98	102	88	107	89	94	88	79	93	89	10	1	1			
	2015	AFRIKA CELIE		5	5	16	25	5	36	24	36	26	27	29	33		1				
CHARMEUR FROM SOUS AFRICA (POETIN)		SEMARGL ADZHEY	3.20	110	101	85	90	92	106	101	87	95	93	87	89	2			2	2	1
	2016	FROM SOUS AFRIKA BELLA TEMBI 2		11	26	5	12	5	26	11	25	13	15	17	21						1
AFRIKA LEONARDO III		LINGA LONGA HARLEY 3	6.35	140	140	95	101	105	95	107	120	121	112	107	114	31	5	3			
	2019	AFRIKA SONJA 4		38	56	29	45	20	58	44	57	46	47	49	53		3				
IMMER MOED LAU		GIANTS OF AFRICA MATTEO	5.41	124	100	94	92	92	112	90	96	96	92	99	96	4		1	4	4	1
	2019	IMMER MOED USHI		5	23	5	14	5	29	13	28	15	17	20	25		1				1

Estimated Breeding Value Indices (EBVi's)

Dams

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-r ament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
IMMER MOED USHI	2014	IMMER MOED OKKE	29.99	121	94	96	80	81	120	78	85	78	73	79	79	2			2	2	1
		LINJO ROXY		5	5	5	5	5	23	5	21	5	5	5	6						1
IMMERMOED NOSTRA DAME	2016	DELATRO GREGOR-KLEINTJIE	9.85	110	98	94		91	102	99	89	94	97	87	90	1			1	1	
		IMMERMOED ANNA BELLA		5	19	5		5	27	5	26	7	12	16	22						
BREKER HURRICANE HATTIE	2017	MAKULU LAANIE ROEPER	9.99	106	103	80	89	80	103	80	100	87	81	92	89	2		1	2	2	1
		BREKER NAOMI 2		5	5	9	39	7	50	39	50	40	42	43	47						1
ALEWIJN'S IMMERMOED AMMEHOELA	2019	CHARMEUR FROM SOUS AFRICA (POETIN)	8.05	116	98	90	95	91	104	99	87	94	94	88	89	1		1			
		IMMERMOED NOSTRA DAME		28	51	5	5	5	15	5	13	5	5	5	5						
ALEWIJNS IMMERMOED BOMBARIE	2020	IMMER MOED LAU	3.85	115	102	87	90	87	108	86	99	93	86	95	93	4	1	1			
		BREKER HURRICANE HATTIE		5	13	5	17	5	30	16	30	17	20	22	26						

