

Boerboel Genetic Evaluation

SA Stud Book
February 2026

MR I.P. MOGOERANE

10410 ZONE 4

0

THABA NCHU

9780



Kennel Number: 728253



Estimated Breeding Values (EBVs)

Litter size and conformation traits are influenced by many genes, with every gene making a small contribution to the expression of the trait. These traits are, however, also influenced by physiological and physical environmental factors, for example the age and sex of the animal, the feeding and management the animal received and year x season effects. It is therefore not possible to know what the animal's genetic potential for the trait is by merely using the measurement.

With BLUP (*Best Linear Unbiased Prediction*) methodology an animal's genetic potential is predicted by making use of the genetic parameters (heritabilities and genetic correlations) of the breed, pedigree information, as well as the measurements of the animal as deviation from the average of the animals that were measured with the animal. Adjustments are made for differences in sex, age, classifier, management and year x season effects to ensure that fair comparisons are made amongst the animals. These predictions are called **EBVs (Estimated Breeding Values)** which indicates the animal's genetic potential for the specific trait and predicts the type of progeny that the animal will breed.

Pedigree information determines the genetic level at which the animal enters the genetic evaluation, it links the information of all relatives together and it is used to determine the genetic ranking of the breed. For young animals that were not yet measured, mid-parent EBVs are estimated, which are the average of the EBVs of the sire and dam. These breeding values are still inaccurate and can change a lot when more information (measurements) is included in subsequent genetic evaluations.

Breeding values therefore enables a breeder to compare his/her animals genetically to one another, as well as to all other animals of the breed. This Report gives a summary of the genetic levels of the active males and females of the relevant breeder in comparison to that of the breed for linearly classified traits and litter size. The genetic change that occurred over time can also easily be seen.

Breeding Value Indices (EBVi)

For easy interpretation, EBVs are standardised and expressed relative to the average (100) of the active animals of the breed, using 12 as standard deviation (SD). This means that an EBV of 100 is equal to the active average of the breed, and more or less 50% of the animals will have EBVi's of between 90 and 110, which indicates an average animal. Only 0.5% of the EBVi's will be lower than 64 and 0.5% will be higher than 136. Higher EBVi's indicate better genetic potential for the trait.

Accuracy

EBVs are published with an **accuracy value (%)**, which indicates how much info was available for the estimation of the EBV and with how much trust it can therefore be applied. Factors that influence the accuracy include heritability, completeness and correctness of pedigree information, number of own and measured relatives and the size of the group in which the animal was measured. When an accuracy of 85% and more is reached, the EBV should not change significantly with the inclusion of more information in subsequent genetic evaluations.

Kennel and Breed Averages: The different traits' averages of the breeder's active animals are compared to those of the breed in the Kennel and Breed Average Graphs and Tables. The breeder can therefore easily see for which traits the animals are above or below the average of the breed.




Genetic Trends: The Genetic Trends indicate the genetic change that occurred over time in the animals bred by the breeder compared to that of the breed. These are calculated as the average of the breeding values of all measured animals bred by the breeder, per year of birth.

Inbreeding: Inbreeding is the mating of related animals and is widely viewed as being harmful. The danger of inbreeding lies there in that it can gradually decrease fertility and survivability – a phenomenon known as inbreeding depression. It can cause lower conception rates, more abortions, more stillborn and weak offspring and a higher susceptibility to diseases. Indications in dogs include a low litter size as well as problems relative to fertility, such as invisible heat, behavioural problems when mating and low sperm quality. Overall, reduced litter size most likely reflects inbreeding depression in several traits, both physiological and behavioural, as well as a high frequency of non-viable embryos due to lethal alleles. Generally, the effects of inbreeding can become noticeable at a value of 6.25%, therefore it is generally recommended that individuals should have inbreeding coefficients of less than 6.25%. Breeders should therefore avoid matings causing inbreeding of more than 6.25% in the progeny. The theoretical maximum inbreeding coefficient is 50%. However, it should be noted that not all inbred animals show signs of inbreeding depression.

The data used in this evaluation was obtained from LOGIX from linear classifications that were done by the Breeder Society. All possible attempts have been made to present the information correctly. SA Stud Book does not accept responsibility for the use and interpretation of the information presented on animals in this publication.

Genetic Levels

Kennel Average relative to Breed Average, where Breed Average is indicated as 0



Average Breeding Value Indices (EBVi) where Breed Average = 100

Trait	All Animals	Males	Females
Litter size (LS)	85	93	81
Movement (Mv)	102	101	102
Temperament (Tmp)	99	98	99
Coat (Ct)	89	91	88
General Appearance	103	104	103
Head & Face (H&F)	104	106	103
Front Quarter (FQ)	93	97	91
Centre (C)	105	102	106
Hind Quarter (HQ)	102	97	105
Final Score (FS)	102	101	102
Hip Health (HH)	92	96	91
Elbow Health (ELM)	98	102	97

Inbreeding

Notes

Inbreeding primarily leads to decreased fertility, low survivability of pups and low disease resistance.

Inbreeding coefficients are calculated for all animals of the breed and indicates the probability that two genes of a pair are the same due to a common ancestor in the pedigree of the animal.

The adverse effects of inbreeding can be observed from 6.25% onwards.


It is therefore highly recommended to record pedigree information as accurately. Inbreeding is dependent on pedigree information. Conventional methods for the calculation of inbreeding coefficients usually assume that unknown parents are unrelated, which is usually far from the truth. Inbreeding coefficients are therefore conservative indicators of the level of inbreeding.

Incomplete or incorrect pedigree information can be reported to SA Stud Book via the Breed Society.

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Rate of Inbreeding



Inbreeding Levels

	All Animals		Males		Females		Sires		Dams	
	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed
Number	11	49007	3	25141	8	23857	5	3371	4	6287
Average Inbreeding	7.77	8.49	6.57	8.46	8.22	8.53	7.64	8.01	8.60	8.00
Number inbred	11	48727	3	24990	8	23728	5	24990	4	23728
Number >6.25% Inbr.	9	30600	1	15537	8	15058	3	15537	2	15058

Estimated Breeding Value Indices (EBVi's)

Males

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	litters
PAURON DUTCH		VILLA ROUGE GOLIAT	8.73	88	104	92	95	88	95	102	94	86	101	94	93	26	5	3			1
	2020	DELMA DONNA		5	5	25	49	22	65	49	64	50	52	55	60						1
PAURON MUFASA		PAURON ROEPER	5.85	102	98	84	105	103	90	116	106	99	107	94	102						
	2021	DELMA DONNA		5	5	12	42	12	60	41	59	43	45	48	54						
PAURON BEES		KILIAN BEES	5.12	99	104	104	104	103	89	94	118	106	98	102	107						
	2022	BREKER LADY		5	5	20	45	20	62	45	61	46	48	51	56						

Estimated Breeding Value Indices (EBVi's)

Females

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information

Inbreed

Health

Fertility

General

Appearance

Progeny :

Appraisal

Health

Progeny with litters

Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Progeny	Litters Litters	Progeny Kennels	Hips Elbows	Progeny with litters
BOSTU LUWELA	2022	KAMMO BBD HONDO	8.39	93	92	73	108	108	83	104	111	95	110	113	109	7	2			
		BOSTU ZHARINA		5	5	36	48	18	64	47	63	48	51	53	58					
BLACKMOUNTAIN NALEDI	2024	PAURON DUTCH	8.19	91	98	83	101	98	89	103	102	90	106	103	101					
		BOSTU LUWELA		5	5	8	26	5	40	26	39	27	29	32	36					
BLACKMOUNTAIN ZAHRA	2024	PAURON DUTCH	8.19	91	98	83	101	98	89	103	102	90	106	103	101					
		BOSTU LUWELA		5	5	8	26	5	40	26	39	27	29	32	36					
BLACKMOUNTAIN IMANI	2025	PAURON DUTCH	8.19	91	98	83	101	98	89	103	102	90	106	103	101					
		BOSTU LUWELA		5	5	8	26	5	40	26	39	27	29	32	36					
BLACKMOUNTAIN ZURI	2025	PAURON DUTCH	8.19	91	98	83	101	98	89	103	102	90	106	103	101					
		BOSTU LUWELA		5	5	8	26	5	40	26	39	27	29	32	36					
BLACKMOUNTAIN AZANA	2025	PAURON DUTCH	8.19	91	98	83	101	98	89	103	102	90	106	103	101					
		BOSTU LUWELA		5	5	8	26	5	40	26	39	27	29	32	36					
BLACKMOUNTAIN AKILI	2025	PAURON DUTCH	8.19	91	98	83	101	98	89	103	102	90	106	103	101					
		BOSTU LUWELA		5	5	8	26	5	40	26	39	27	29	32	36					
BLACKMOUNTAIN AYANA	2025	PAURON DUTCH	8.19	91	98	83	101	98	89	103	102	90	106	103	101					
		BOSTU LUWELA		5	5	8	26	5	40	26	39	27	29	32	36					

Estimated Breeding Value Indices (EBVi's)

Sires

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Sire Information		Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels	Hips	Elbows	
VILLA ROUGE GOLIAT	2013	AUGUSTYN BACCHUS	12.33	83	112	103	97	79	104	77	87	81	92	86	85	143	18	16		14
		BUNDU VALLEY BBD ELSA		5	5	49	66	41	79	66	78	67	69	71	75		13			4
KILIAN BEES	2018	MES BITE A BULLET	7.09	82	83	111	111	104	95	113	115	99	108	108	111	2	1			
		KLEIN SANDFONTEIN GRACI		5	17	41	55	35	69	55	68	56	58	60	64		1			
KAMMO BBD HONDO	2018	MIDDELPOS BBD MAXIMUS I	5.64	108	92	99	110	102	101	113	108	94	101	98	104	58	8	15		4
		AREA BOERBOELS KARLIEN VAN JAARVELD		5	14	41	60	37	74	60	73	61	63	65	69		11			2
PAURON ROEPER	2019	MAKULU LAANIE ROEPER	4.38	111	100	87	97	96	87	102	101	96	102	91	96	27	4	3		2
		SAMRICO BEKKA		5	5	22	54	27	71	53	70	55	57	60	65		3			1
PAURON DUTCH	2020	VILLA ROUGE GOLIAT	8.73	88	104	92	95	88	95	102	94	86	101	94	93	26	5	3		1
		DELMA DONNA		5	5	25	49	22	65	49	64	50	52	55	60		3			1

Estimated Breeding Value Indices (EBVi's)

Dams

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name		Sire	%	Hips	Elbows	Litter Size	Move-	Tempe-	Coat	General	Head &	Front	Centre	Hind	Final	Progeny	Litters	Progeny Kennels	Hips	Elbows	
	BYr	Dam		Acc.	Acc.	Acc.	ment	rament		Appear.	Face	Quarter		Quarter	Score						
DELMA DONNA		MES RAMBO 1	5.16	93	96	80	111	110	81	124	111	103	117	105	110	11	1	5		3	
	2017	ELEVATION LOLA		5	5	40	56	33	71	55	70	56	59	61	66			4	1		
BREKER LADY		MIDDELPOS BBD GAMKA	3.87	117	124	97	85	96	75	83	114	108	88	94	97	12	4	4		1	
	2020	BREKER NELLA		5	15	36	60	38	73	59	72	61	62	65	69			3	1		
BOSTU ZHARINA		BOSTU ZAZU	17.00	78	92	52	103	108	84	99	109	93	110	117	109	15	3	7		3	
	2021	BOSTU XHENA		5	5	50	59	30	75	58	74	60	63	65	70			5	1		
BOSTU LUWELA		KAMMO BBD HONDO	8.39	93	92	73	108	108	83	104	111	95	110	113	109	7	2				
	2022	BOSTU ZHARINA		5	5	36	48	18	64	47	63	48	51	53	58						

