

Boerboel Genetic Evaluation

SA Stud Book
February 2026

MR I.M. NDLOVU
THORNVILLE AMBLETON 2
0
PIETERMARITZBURG
4004



Kennel Number: 717805



Estimated Breeding Values (EBVs)

Litter size and conformation traits are influenced by many genes, with every gene making a small contribution to the expression of the trait. These traits are, however, also influenced by physiological and physical environmental factors, for example the age and sex of the animal, the feeding and management the animal received and year x season effects. It is therefore not possible to know what the animal's genetic potential for the trait is by merely using the measurement.

With BLUP (*Best Linear Unbiased Prediction*) methodology an animal's genetic potential is predicted by making use of the genetic parameters (heritabilities and genetic correlations) of the breed, pedigree information, as well as the measurements of the animal as deviation from the average of the animals that were measured with the animal. Adjustments are made for differences in sex, age, classifier, management and year x season effects to ensure that fair comparisons are made amongst the animals. These predictions are called **EBVs (Estimated Breeding Values)** which indicates the animal's genetic potential for the specific trait and predicts the type of progeny that the animal will breed.

Pedigree information determines the genetic level at which the animal enters the genetic evaluation, it links the information of all relatives together and it is used to determine the genetic ranking of the breed. For young animals that were not yet measured, mid-parent EBVs are estimated, which are the average of the EBVs of the sire and dam. These breeding values are still inaccurate and can change a lot when more information (measurements) is included in subsequent genetic evaluations.

Breeding values therefore enables a breeder to compare his/her animals genetically to one another, as well as to all other animals of the breed. This Report gives a summary of the genetic levels of the active males and females of the relevant breeder in comparison to that of the breed for linearly classified traits and litter size. The genetic change that occurred over time can also easily be seen.

Breeding Value Indices (EBVi)

For easy interpretation, EBVs are standardised and expressed relative to the average (100) of the active animals of the breed, using 12 as standard deviation (SD). This means that an EBV of 100 is equal to the active average of the breed, and more or less 50% of the animals will have EBVi's of between 90 and 110, which indicates an average animal. Only 0.5% of the EBVi's will be lower than 64 and 0.5% will be higher than 136. Higher EBVi's indicate better genetic potential for the trait.

Accuracy

EBVs are published with an **accuracy value (%)**, which indicates how much info was available for the estimation of the EBV and with how much trust it can therefore be applied. Factors that influence the accuracy include heritability, completeness and correctness of pedigree information, number of own and measured relatives and the size of the group in which the animal was measured. When an accuracy of 85% and more is reached, the EBV should not change significantly with the inclusion of more information in subsequent genetic evaluations.

Kennel and Breed Averages: The different traits' averages of the breeder's active animals are compared to those of the breed in the Kennel and Breed Average Graphs and Tables. The breeder can therefore easily see for which traits the animals are above or below the average of the breed.




Genetic Trends: The Genetic Trends indicate the genetic change that occurred over time in the animals bred by the breeder compared to that of the breed. These are calculated as the average of the breeding values of all measured animals bred by the breeder, per year of birth.

Inbreeding: Inbreeding is the mating of related animals and is widely viewed as being harmful. The danger of inbreeding lies there in that it can gradually decrease fertility and survivability – a phenomenon known as inbreeding depression. It can cause lower conception rates, more abortions, more stillborn and weak offspring and a higher susceptibility to diseases. Indications in dogs include a low litter size as well as problems relative to fertility, such as invisible heat, behavioural problems when mating and low sperm quality. Overall, reduced litter size most likely reflects inbreeding depression in several traits, both physiological and behavioural, as well as a high frequency of non-viable embryos due to lethal alleles. Generally, the effects of inbreeding can become noticeable at a value of 6.25%, therefore it is generally recommended that individuals should have inbreeding coefficients of less than 6.25%. Breeders should therefore avoid matings causing inbreeding of more than 6.25% in the progeny. The theoretical maximum inbreeding coefficient is 50%. However, it should be noted that not all inbred animals show signs of inbreeding depression.

The data used in this evaluation was obtained from LOGIX from linear classifications that were done by the Breeder Society. All possible attempts have been made to present the information correctly. SA Stud Book does not accept responsibility for the use and interpretation of the information presented on animals in this publication.

Genetic Levels

Kennel Average relative to Breed Average, where Breed Average is indicated as 0



Average Breeding Value Indices (EBVi) where Breed Average = 100

Trait	All Animals	Males	Females
Litter size (LS)	105	106	104
Movement (Mv)	98	104	91
Temperament (Tmp)	102	106	98
Coat (Ct)	99	99	100
General Appearance	104	104	104
Head & Face (H&F)	104	105	103
Front Quarter (FQ)	103	105	101
Centre (C)	102	102	102
Hind Quarter (HQ)	102	106	99
Final Score (FS)	103	105	100
Hip Health (HH)	100	102	97
Elbow Health (ELM)	102	104	99

Inbreeding

Notes

Inbreeding primarily leads to decreased fertility, low survivability of pups and low disease resistance.

Inbreeding coefficients are calculated for all animals of the breed and indicates the probability that two genes of a pair are the same due to a common ancestor in the pedigree of the animal.

The adverse effects of inbreeding can be observed from 6.25% onwards.

It is therefore highly recommended to record pedigree information as accurately. Inbreeding is dependent on pedigree information. Conventional methods for the calculation of inbreeding coefficients usually assume that unknown parents are unrelated, which is usually far from the truth. Inbreeding coefficients are therefore conservative indicators of the level of inbreeding.

Incomplete or incorrect pedigree information can be reported to SA Stud Book via the Breed Society.

Rate of Inbreeding

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Inbreeding Levels

	All Animals		Males		Females		Sires		Dams	
	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed
Number	4	49007	2	25141	2	23857	4	3371	4	6287
Average Inbreeding	7.72	8.49	6.88	8.46	8.55	8.53	8.25	8.01	6.81	8.00
Number inbred	4	48727	2	24990	2	23728	4	24990	4	23728
Number >6.25% Inbr.	3	30600	1	15537	2	15058	3	15537	2	15058

Estimated Breeding Value Indices (EBVi's)

Males

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
SAFIKA TORNADO		KYLANO TAXMAN	5.93	103	105	106	126		112	104	95	108	106	111	108						
	2022	HEAVY WEIGHTS ZANDA		5	5	13	41		58	40	57	42	44	47	52						
ALPHUS MKHONTO		MES DIMO	7.84	100	103	107	83	98	86	105	115	102	99	101	103						
	2024	SAFIKA SANDY		5	5	5	26	5	38	26	37	27	29	31	34						

Estimated Breeding Value Indices (EBVi's)

Females

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe- rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels		Hips	Elbows	
SAFIKA HUSKY		HEAVY WEIGHTS VOLCANO	7.52	98	94	103	86	102	107	116	111	108	101	92	104						
	2022	SAFIKA CLIVIA		5	5	5	38	5	55	37	54	39	41	44	49						
SHANSAFRICA VOLCANIC VICKY		BOUWER MAXIMUS 2	9.59	97	104	106	97	94	93	92	94	94	102	106	97						
	2024	REYDALL HANNAH		5	5	5	16	5	35	15	34	17	20	24	29						

Estimated Breeding Value Indices (EBVi's)

Sires

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Sire Information			Inbreed	Health		Fertility	General			Appearance						Progeny :	Appraisal	Health	Progeny
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels	Hips Elbows	with litters
KYLANO TAXMAN	2018	CRIHATA TROY	6.46	107	105	94	121	115	118	105	102	109	101	104	108	49 14	13 10		3 1
		KAMMO BBD ASTRID		5	5	20	54	24	72	54	71	55	58	61	66				
MES DIMO	2019	MES MAC GYVER	4.70	102	111	111	104	108	82	97	108	99	100	111	105	12 3	1 1		
		CABARET MANDIE		5	5	33	49	27	59	48	59	49	51	53	56				
HEAVY WEIGHTS VOLCANO	2021	AFRIKA GHUNTER	10.81	92	82	94	70	106	68	119	121	106	108	99	106	19 2	8 7		3 1
		JOSMOT MISS S.A.		5	5	24	59	33	74	59	73	60	62	65	69				
BOUWER MAXIMUS 2	2021	KAMMO BBD BOTTER	11.03	93	108	103	90	95	91	93	101	97	109	116	104	8 1			
		BOUWER PRADA		5	5	5	35	5	51	34	50	36	38	41	46				

Estimated Breeding Value Indices (EBVi's)

Dams

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe- r ament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
HEAVY WEIGHTS ZANDA	2018	BOUWER STOFFEL 1	8.70	100	106	117	115	105	117	107	97	112	106	103	106	19	4	9			3
		BOUWER SASHA 1		5	5	41	49	17	67	49	66	50	53	56	61			7			1
SAFIKA CLIVIA	2020	KYLANO TAXMAN	5.93	103	105	113	117	105	134	110	92	106	100	101	104	23	3	8			1
		HEAVY WEIGHTS ZANDA		5	5	19	47	14	66	46	65	48	51	54	60			6			1
SAFIKA SANDY	2022	HEAVY WEIGHTS VOLCANO	5.31	98	95	102	61	88	90	113	121	105	97	91	101	5	2				
		SUBIRI KHLOE		5	5	20	41	11	57	41	56	42	44	47	51						
REYDALL HANNAH	2022	REYDALL PIET	7.30	101	100	109	103	94	94	90	87	90	95	96	90	8	1				
		REYDALL NIXIE		5	5	5	37	5	56	36	54	38	40	43	49						

