



DOHNE MERINO GENETIC REPORT



20 September 2026

SOUTHGATE

FDM : AC



CERTIFICATE
OF QUALITY

Valid up to
October 2029



Information

This Report gives a summary of the genetic levels of the flock with regards to production and fertility. Breeders can easily see if genetic progress occurred in their flocks.

EBV (Estimated Breeding Value)

The breeding value indicates an animal's genetic ability for a specific trait. It is indicated in the unit of measurement and relative to that of an average animal born in the base year, i.e. five years ago. A higher breeding value is desirable for all traits, except Fibre Diameter (thinner fibres are better), AFL (early sexual maturity) and ILP (easy conceptions – shorter Interlambing Periods).

EBVi (EBVindex)

The base of a breeding value index is 100, where 100 is the average of the active animals in the breed. Values above 100 are always in the more desirable direction, also for AFL (early sexual maturity) and ILP (shorter intervals). However, for FD values below 100 indicate thinner (more desirable) fibres.

Selection Indices

This is the combination of breeding values into a single value. It indicates of the genetic value of the combined traits of the animal with regards to the breed. Traits cannot simply be summed, as they are expressed in different units, their genetic variation differ and because negative values are more desirable for certain traits. Traits are also weighted differently according to economic importance. All these factors are taken into account when selection indices are constructed. The selection indices currently available for the Dohne Merino breed, is the General Merit (GM) and the within-test-group selection index (SIP). The GM and SIP indices are only published for animals with wean and wool measurements.

General Merit (GM)

$$GM = (EBV \text{ BWeight} * 2.30) + (EBV \text{ CFW} * 22.0) + (EBV \text{ FD} * -5.0) + (EBV \text{ TWW} * 5.0)$$

Within-Test-Group Selection Index (SIP)

$$SIP = (GM_{dier} / \text{Average GM of wool contemporary group}) * 100$$

This is a lifetime value for the animal, allocated on the basis of the animal's own measurements. This value is not updated when the animal's EBVs are estimated more accurately due to the inclusion of progeny and other relatives' information in the Genetic Evaluation.

Genetic Windows and Averages

The flock's average genetic levels are compared to the active breed average, where active breed average is set to 100.

Genetic Trends

The Genetic Trends indicate the genetic changes that occurred in the flock over time in comparison to that of the breed. This is based only on the animals that were measured for the relevant trait.

Inbreeding

The average inbreeding of the flock per year of birth is compared to that of the breed. Inbreeding levels should stay below 6.25% and rate of inbreeding less than 0.1-0.2% per year to maintain genetic diversity.

The active population is defined as the lambs born in the last 36 months and their parents, regardless if they were measured for the trait or not, excluding the animals that are already cancelled on Logix.

Abbreviations

WnD:	Weaning Weight Direct
WnM	Weaning Weight Maternal
BW:	Body Weight
CFW:	Clean Fleece Weight
FD:	Fibre Diameter
SL:	Staple Length
AFL:	Age First Lambing
ILP:	Interlambing Period
NLW:	Number of Lambs Weaned
TWW:	Total Weight of Lambs Weaned

The data used in this evaluation has been obtained from LOGIX, as supplied by breeders. Everything possible had been done to present the information correctly. SA Stud Book does not take any responsibility for the use and interpretation of the information from this publication.

Genetic Windows

Average Breeding Value Indices (EBVi)

(Lambs born in the last 36 months and their parents)

Number: 187


Genetic Linkage

(Based on the youngest measured generation)

Level	Nr of Link Rams
Geen / None	0

A link ram should have at least 20 measured progeny in your flock and at least another 20 measured progeny in at least one more flock.

Try to use at least 5 link rams in a 5-year period to obtain strong genetic linkages within the breed.



Abbr.	Trait	Interpretation of Index	
		<100 (Worse)	>100 (Better)
WnD	Weaning Weight Direct	Lighter	Heavier
WnM	Weaning Weight Maternal	Less	More
BW	Body Weight	Lighter	Heavier
CFW	Clean Fleece Weight	Less	More
FD	Fibre Diameter	Thinner	Thicker
SL	Staple Length	Shorter	Longer
AFL	Age at First Lambing	Later	Earlier
ILP	Interlambing Period	Longer	Shorter
NLW	Number of Lambs Weaned	Less	More
TWW	Total Weight of Lambs Weaned	Lighter	Heavier
AM	General Merit	Poor	Good
SIP	Within-Test Group Selection Index	Poor	Good

* The Genetic Windows provide a quick overview of the flock's genetic status compared to the breed.

* The values presented here are the average breeding value indices and selection indices of all the lambs born in the past 36 months and their parents in the flock compared to the breed.


* An index of 100 equals the average of the lambs born in the past 36 months and their parents in the breed.

* The breeding value indices for AFL and ILP are expressed in such a way that an index above 100 indicates the better animals, thus the animals with a genetic ability for a short AFL and ILP. See accompanying table for interpretation of other Indices.

Genetic Windows


Sires of lambs born in the last 36 months

Number : 7




Dams of lambs born in the last 36 months

Number : 53



Lambs born in the last 36 months

Number : 90



Average Genetic Levels

Average Breeding Values

Averages of all lambs born in the last 36 months and their parents

Trait	Flock			Breed		
	All	Rams	Ewes	All	Rams	Ewe
Breeding Values for Production Traits						
Weaning Weight Direct	-0.24	-0.02	-0.37	0.54	0.66	0.47
Weaning Weight Mat	0.05	0.02	0.06	0.01	0.00	0.01
Body Weight	-0.13	0.21	-0.34	0.56	0.96	0.41
Clean Fleece Weight	-0.11	-0.09	-0.12	0.03	0.06	0.02
Fibre Diameter	0.13	0.05	0.18	-0.14	-0.22	-0.12
Staple Length	1.56	2.81	0.78	0.50	0.68	0.43
Breeding Values for Fertility						
Age First Lambing	-7.55	-7.08	-7.87	1.48	2.04	1.17
Interlambing Period	-2.36	-2.20	-2.47	0.36	0.44	0.31
Lambs Weaned	1.61	1.81	1.49	0.46	0.56	0.39
Total Weight Weaned	0.24	0.28	0.22	0.27	0.32	0.24
Selection Indices						
General Merit Index	-1.72	0.26	-2.71	2.81	4.45	2.23
SIP	99.9	100.6	99.5	100.7	100.4	100.8
Inbreeding						
Inbreeding (%)	2.31	2.52	2.17	3.00	3.13	2.92
Number of Animals						
Number of Animals	189	71	118	125905	45967	79938









Average Breeding Value Indices

Average EBVi of all active measured animals, where breed average of active measured animals = 100.

Trait	Flock		
	All	Rams	Ewes
Breeding value Indices for Production Traits			
Weaning Weight Direct	92.7	94.7	91.4
Weaning Weight Mat	100.8	100.3	101.2
Body Weight	95.6	97.8	94.2
Clean Fleece Weight	91.2	92.7	90.4
Fibre Diameter	104.1	102.9	104.9
Staple Length	102.8	106.1	100.7
Breeding Value Indices for Fertility			
Age First Lambing	110.2	109.6	110.5
Interlambing Period	109.1	108.5	109.4
Lambs Weaned	104.3	105.1	103.9
Total Weight Weaned	99.7	100.1	99.4
Selectopn Indices			
General Merit Index	92.5	95.8	90.9
SIP	99.9	100.6	99.5
Inbreeding			
Inbreeding (%)	2.31	2.52	2.17
Number of Animals			
Number of Animals	189	71	118

Genetic Trends




Average breeding value per year of birth for animals in the breed and flock. Genetic trends are based on measured animals that were bred by the relevant breeder.



For FD, AFL and ILP, flock trends below breed averages are desirable

Genetic Trends: Fertility Traits

Average breeding value per year of birth in the breed and flock. Genetic trends are based on measured animals that were bred by the relevant breeder.



Genetic Trends: Breed and Flock

Average breeding value per year of birth for animals in the breed and flock. Genetic trends are based on measured animals that were bred by the relevant breeder.

Genetic Trends of Breed


Birth Year	Weaning Weight			Body Weight		CFW		FD		SL		AFL		ILP		NLW		TWW		General Merit	
	Nr	Dir	Mat	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	Ind
2014	34600	-1.62	-0.08	23613	-2.43	23477	-0.20	23511	0.33	23404	-1.65	6513	-4.96	4764	-1.69	8149	-0.57	6758	-0.41	25195	-9.76
2015	36007	-1.44	-0.05	25062	-2.12	24975	-0.18	24990	0.28	24511	-1.51	6856	-4.79	5313	-1.43	8627	-0.57	7110	-0.42	26207	-8.59
2016	33764	-1.17	-0.03	24799	-1.74	24614	-0.15	24664	0.22	24080	-1.07	7054	-4.46	5449	-1.38	8681	-0.41	7293	-0.27	25836	-6.81
2017	33408	-0.95	0.00	24007	-1.39	23924	-0.13	23948	0.19	23487	-1.19	7125	-3.04	5498	-0.77	9139	-0.30	7020	-0.19	24328	-5.48
2018	34871	-0.67	0.00	24937	-0.96	24789	-0.10	24807	0.17	24087	-0.72	7421	-2.86	5764	-0.67	8663	0.09	7374	0.04	25402	-3.35
2019	34833	-0.46	0.02	25157	-0.74	25034	-0.06	25045	0.07	24985	-0.63	7016	-1.62	5333	-0.50	8420	-0.19	6949	-0.05	25370	-2.61
2020	35032	-0.26	0.03	24234	-0.34	24298	-0.04	24310	0.04	24213	-0.20	6557	-0.74	4885	-0.15	8732	0.34	6476	0.18	24153	-0.71
2021	33757	0.00	0.00	23306	0.00	23200	0.00	23217	0.00	23156	0.00	6425	0.00	4779	0.00	8439	0.00	6245	0.00	23180	0.17
2022	32965	0.20	-0.02	22207	0.34	22115	0.01	22136	-0.05	22041	0.27	5933	0.83	3883	0.50	7685	0.48	5658	0.21	22071	1.75
2023	32993	0.44	-0.06	21947	0.60	21831	0.04	21880	-0.16	21853	0.46	5361	1.77	1851	1.14	5554	0.35	4743	0.22	21836	2.88
2024	34418	0.50	-0.04	22757	0.76	22780	0.06	22790	-0.16	22746	0.74	2626	4.31	147	0.21	2149	0.65	2008	0.48	22746	3.82
2025	28497	0.77	0.02	8675	1.57	9028	0.07	9040	-0.38	9011	0.95	60	10.56							8597	6.80

Genetic Trends of Flock

Birth Year	Weaning Weight			Body Weight		CFW		FD		SL		AFL		ILP		NLW		TWW		General Merit	
	Nr	Dir	Mat	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Nr	EBV	Aantal	EBV	Nr	EBV
2014	46	-3.05	0.09	9	-5.20	9	-0.34	9	0.88	9	-0.92									9	-17.60
2015	19	-2.70	0.30	15	-4.22	15	-0.29	15	0.60	15	-2.64	8	-6.84	7	-2.57	15	1.42	8	0.05	15	-13.10
2016	32	-2.64	-0.20									20	-17.86	14	-4.87	20	-0.71	20	-0.93	31	-14.39
2017	43	-1.91	-0.09	36	-3.40	36	-0.20	36	0.47	36	2.20	10	-17.98	8	-3.67	11	1.38	11	-0.11	36	-11.31
2018	63	-2.73	-0.18	58	-4.64	58	-0.21	58	0.58	58	0.06	14	-16.11	13	-4.27	14	-0.76	14	-0.63	58	-15.13
2019	42	-1.67	0.18	24	-1.86	24	-0.30	24	0.45	24	0.21	11	-6.36	8	-1.40	20	1.00	11	0.22	24	-9.63
2020	51	-1.66	-0.12	35	-2.74	35	-0.16	35	0.20	35	-2.35	9	-6.78	4	-2.42	9	0.51	9	0.12	35	-7.08
2021	68	-0.98	-0.03	41	-1.63	41	-0.07	41	0.58	41	-1.15	14	-10.79	9	-4.53	14	1.15	14	-0.19	41	-6.57
2022	54	-0.76	0.04													13	0.45				
2024	42	-0.64	0.06	8	-0.38	8	-0.15	8	-0.41	8	4.27									8	-0.31
2025	66	0.23	0.03	55	0.39	53	-0.04	53	0.28	53	1.66									53	-0.07

Inbreeding

Inbreeding in Flock and Breed



Pedigree Information

	Number	%
Known Sires	177	94
Known Dams	177	94

Please Note : When pedigree information is incomplete, inbreeding is under-estimated.

The effects of inbreeding can be observed from 6.25% onwards. Inbreeding primarily leads to decreased fertility and low survivability of lambs.

Inbreeding % is dependent on the accuracy of pedigree information. It is therefore highly recommended to record pedigrees as accurately and completely as possible and not to mate related animals.

Incomplete and incorrect pedigree information can be reported to SA Stud Book

Inbreeding Level and Rate

Year of Birth	Level* (%)	Rate (% / year)
2000 - 2025	2.15	0.03
Recommendations	< 6.25	< 0.1 - 0.2

*Of last birth year

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Percentiles of Breeding Values : All Active Animals

Percentile	Wean Dir	Wean Mat	Body Weight	CFW	FD	SL	NLW	TWW	AFL	ILP	Gen. Merit
Gemid/Avg	0.43	0.00	0.58	0.04	-0.15	0.46	0.47	0.26	1.30	0.30	2.53
SA/SD	1.29	0.56	1.84	0.17	0.69	3.92	2.99	0.93	-10.19	-3.48	7.11
MIN	-4.90	-3.44	-9.56	-0.75	3.11	-18.57	-17.30	-5.29	68.72	22.49	-25.62
1%	-2.32	-1.35	-3.42	-0.38	1.49	-8.80	-7.64	-2.17	33.49	9.02	-13.42
5%	-1.58	-0.89	-2.30	-0.25	0.95	-5.86	-4.59	-1.30	18.37	6.15	-8.27
10%	-1.19	-0.67	-1.72	-0.19	0.69	-4.38	-3.19	-0.87	13.14	4.69	-6.12
20%	-0.68	-0.43	-0.98	-0.11	0.41	-2.72	-1.72	-0.43	8.52	3.05	-3.50
30%	-0.30	-0.27	-0.43	-0.05	0.20	-1.54	-0.76	-0.11	5.14	1.87	-1.46
40%	0.05	-0.13	0.05	0.00	0.02	-0.54	0.06	0.10	2.16	0.90	0.38
50%	0.38	0.00	0.50	0.04	-0.14	0.39	0.45	0.21	0.19	0.13	1.99
60%	0.72	0.13	0.96	0.08	-0.31	1.32	1.08	0.41	-1.46	-0.57	3.85
70%	1.09	0.27	1.48	0.13	-0.49	2.34	1.79	0.65	-3.58	-1.45	6.03
80%	1.54	0.44	2.11	0.18	-0.72	3.60	2.66	0.96	-5.70	-2.39	8.71
90%	2.13	0.69	2.98	0.26	-1.03	5.43	4.08	1.41	-9.18	-3.66	12.22
95%	2.62	0.93	3.73	0.32	-1.29	7.03	5.33	1.81	-12.92	-4.88	14.90
99%	3.57	1.41	5.08	0.45	-1.80	10.22	7.78	2.59	-23.99	-8.36	19.40
MAX	5.84	2.39	8.39	0.84	-3.43	19.74	16.55	5.29	-43.24	-17.72	31.42

SD = Standard Deviation = Indicates the spread around the average

Percentiles of Indices : All Active Animals

Percentile	Wean Dir	Wean Mat	Body Weight	CFW	FD	SL	NLW	TWW	AFB	ILP	Gen. Merit
Gemid/Avg	100	100	100	100	100	100	100	100	100	100	100
SA/SD	12	12	12	12	12	12	12	12	12	12	12
MIN	49	24	35	52	149	50	33	33	24	30	53
1%	73	71	73	72	128	73	68	69	63	70	72
5%	81	81	81	80	120	80	80	80	80	80	80
10%	85	85	85	85	115	85	85	85	86	85	85
20%	90	91	90	90	110	90	91	91	92	91	90
30%	93	94	94	94	106	94	95	95	95	95	94
40%	97	97	97	97	103	97	98	97	99	98	97
50%	100	100	100	100	100	100	100	100	101	100	100
60%	103	103	103	103	97	103	103	103	103	103	103
70%	106	106	106	106	94	106	106	106	106	106	106
80%	110	109	110	110	90	110	109	109	108	109	110
90%	116	115	115	115	85	115	115	115	112	114	116
95%	120	120	120	120	80	120	119	120	117	118	120
99%	129	130	130	128	72	129	129	129	130	130	128
MAX	150	150	150	150	49	150	160	160	150	160	147

Rams with Progeny



Genotyped

*Inbreeding in red is high

EBV - Estimated Breeding Value

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding values

Breeding values with accuracy below 5% are not printed

Index: Expresses the EBV around 100 where 100 is set as the average EBV of lambs born in the past 36 months and their parents in the breed

Animal Information		Parents & Progeny					Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility				Rank
Animal ID Regnr	% In- breeding	Sire ID Dam ID	Wean		Wool test		SIP	GM		Weaning Weight		Body Weight	CFW	FD	SL	NLW	TWW	DAM		
			Progeny	Nr	Progeny	Nr		EBV	Index	Dir	Mat							Acc	Index	
			weaned	flocks	tested	flocks			EBV	Index	Acc	Index	Acc	Index	Acc	Index	Acc	Index	Acc	Index
		Active Breed Average :					101	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27			
B 21 0072 92689611	4.87	B 19 0199 B 18 0219	85	2	52	2	106	4.78	103	1.26 94	0.01 107	1.10 103	-0.18 90	-0.61 94	1.52 103	6.05 33	0.98 121	127	6	30
B 22 0106 95089371	2.34	B 20 0284 B 20 0321	4	1	1	1	102	0.91	97	-0.10 73	-0.62 94	0.69 101	0.02 67	0.71 77	2.98 113	2.30 33	0.51 107	115	4	61
OJ 22 0186 95844981	1.56	SD 18 0732 OJ 15 0493	2	1	1	1	105	-2.64	91	0.65 70	0.42 101	0.18 98	-0.12 66	0.01 76	-1.80 102	-1.26 35	-0.31 93	118	5	30
RA 19 0914 85526457	1.27	RA 17 0736 RA 15 0858	348	5	281	5	119	10.99	113	1.84 98	0.27 112	2.30 106	0.13 98	-0.80 99	4.75 111	-0.01 67	0.59 104	121	6	1
RA 21 0826 92532647	3.54	RA 19 0314 RA 16 0808	72	2	58	2	104	-0.83	94	0.49 93	-1.03 100	-0.07 92	0.23 91	-0.42 95	7.66 119	-3.98 42	-1.67 83	108	5	76
RA 21 1048 92533785	3.94	RA 19 0450 RA 17 0033	32	1	6	1	107	4.56	103	-0.92 87	0.25 105	0.03 97	-0.25 74	-1.58 82	9.06 122	5.06 37	0.73 117	91	3	43
RA 22 0028 94744364	3.54	RA 19 0914 RA 19 0701	31	1	21	1	109	6.53	106	1.65 87	0.12 110	2.92 115	0.16 83	0.94 90	4.90 112	1.28 41	0.68 103	131	5	24
XT 19 0966 86412624	1.78	RA 15 0742 XT 15 0578	102	2	73	2	111	0.88	97	-0.81 94	-0.36 61	-1.10 93	0.05 92	-0.34 96	-2.65 97	3.62 41	0.15 112	95	2	8
XT 22 0232 95811386	1.67	RA 18 0736 XT 19 0967	27	2	24	2	105	-0.44	95	2.03 86	0.08 114	1.91 109	-0.06 84	0.34 90	0.71 107	-1.43 39	-0.44 93	117	5	28
XT 22 2277 95812079	1.89	XT 20 3024 XT 19 2988	65	2	22	2	118	4.98	104	1.36 92	0.63 108	1.43 106	-0.09 82	-0.80 89	-0.20 90	2.51 20	0.29 108	111	3	3
XT 22 5209 95811121	1.70	XT 19 8920 XT 18 5854	55	2	23	2	108	3.14	101	-0.65 91	-0.13 89	-1.56 85	0.17 83	-0.17 90	0.33 100	1.25 23	0.63 103	106	4	13
Rams with Progeny Average :							109	3.0	100	0.62	-0.03	0.71	0.01	-0.25	2.48	1.40	0.19			

Young Rams



Genotyped

EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding value

Breeding values with an accuracy of below 5% are not printed

Young rams, born since 2020 with a weaning weight, and wool test, but no progeny, both parents known.

Animal Information		Parents		Rank	Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility			
Animal ID Regnr	% In- breeding	Sire ID	Dam ID		SIP	GM		Weaning Dir	Weight Mat	Body Weight	CFW	FD	SL	NLW	TWW	EPI	DAM Lambed
						EBV	Index	EBV	Acc	Index	EBV	Acc	Index	EBV	Acc	Index	Phenotype
		Active Breed Average :			101	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27		
AC 24 0004 101610400	0.97	RA 21 1048	AC 21 0010	5	99	-1.00	94	-0.35 66	0.03 39	-0.45 100	-0.05 61	-0.32 72	2.19 65	1.84 28	-0.20 27	101	2
AC 24 0009 101610467	2.08	OJ 22 0186	AC 22 0106	7	96	-4.44	88	-0.36 55	0.23 31	-1.77 105	-0.20 53	-0.46 67	-5.86 58	-0.41 18	-0.05 18	96	1
AC 24 0018 101610525	2.08	RA 21 1048	AC 22 0114	1	108	4.88	103	-0.04 61	0.29 34	1.86 106	-0.19 56	-0.82 69	4.54 61	2.75 18	0.47 18	105	2
AC 24 0021 101610566	2.15	B 22 0106	RA 22 0116	2	106	4.33	103	0.33 63	-0.47 36	2.16 90	0.03 60	0.63 70	1.12 63	4.06 25	0.68 25	97	1
AC 24 0023 101610582	3.12	RA 21 1048	RA 22 0226	3	101	0.73	97	0.12 68	0.27 40	1.10 106	-0.10 62	0.29 73	12.29 67	1.59 28	0.40 28	113	2
AC 24 0031 101610681	7.28	RA 21 1048	RA 22 0923	8	95	-4.28	88	-2.12 67	-0.46 38	-2.72 90	-0.12 62	-0.19 71	9.27 65	4.23 26	0.34 26	93	1
AC 24 0040 101610764	4.05	RA 21 1048	AC 19 0024	6	97	-1.98	92	-0.48 67	0.61 43	-0.89 113	-0.38 62	-1.09 71	2.95 65	3.23 31	0.40 30	143	4
AC 24 0042 101610806	3.48	RA 21 1048	AC 18 0056	4	99	-0.69	94	-2.22 67	-0.32 42	-2.32 93	-0.19 62	-1.29 73	7.70 66	4.02 30	0.37 29	102	4
AC 25 0005 109703702	1.61	RA 22 0028	XT 22 0226	6	105	3.53	101	1.01 67	0.36 39	1.48 107	-0.05 64	0.22 76	3.63 69	1.96 23	0.71 23	102	1
AC 25 0006 109703710	2.21	RA 22 0028	XT 21 5191	3	107	5.02	104	0.64 61	0.02 33	1.60 100	0.18 59	0.62 72	-2.84 64	0.87 20	0.45 20	102	1
AC 25 0009 109703744	2.02	B 21 0072	XT 21 1119	20	95	-3.27	90	-0.14 69	0.28 40	-0.64 106	-0.25 66	0.16 76	-1.46 70	4.74 25	0.59 24	94	2
AC 25 0016 109703769	2.21	B 21 0072	XT 21 0168	2	108	5.39	104	0.68 64	0.12 35	1.30 102	-0.07 62	-0.56 73	-2.28 66	3.25 16	0.59 16	109	1
AC 25 0018 109703785	2.21	RA 22 0028	XT 21 0170	17	99	-0.98	94	-0.19 63	0.12 36	0.11 102	-0.01 59	0.79 72	3.77 65	0.87 20	0.45 20	107	1
AC 25 0020 109703793	1.48	RA 22 0028	XT 21 1152	22	95	-3.83	89	0.40 67	-0.16 38	1.52 96	-0.04 65	1.36 76	3.47 69	-0.41 28	-0.29 28	86	3
AC 25 0032 109703827	3.73	RA 22 0028	AC 21 0046	8	104	3.07	100	2.30 67	0.40 41	3.26 108	-0.14 64	0.71 75	-2.65 69	2.41 28	0.65 27	109	2

Young Rams



Genotyped

EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding value

Breeding values with an accuracy of below 5% are not printed

Young rams, born since 2020 with a weaning weight, and wool test, but no progeny, both parents known.

Animal Information		Parents		Rank	Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility		
Animal ID Regnr	% In- breeding	Sire ID	Dam ID		SIP	GM		Weaning Weight Dir	Weight Mat	Body Weight	CFW	FD	SL	NLW	TWW	DAM EPI Lambled
					EBV	Index		Acc	Index	Acc	Acc	Index	Acc	Acc	Index	Phenotype
Active Breed Average :					101	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27	
AC 25 0035 109703835	1.33	RA 22 0028	AC 21 0043	19	96	-3.08	90	0.14 67 96	0.07 41 101	0.16 67 97	0.10 64 104	1.85 75 131	9.31 69 123	2.18 28 107	0.44 28 102	105 2
AC 25 0040 109703868	3.28	RA 21 0826	AC 21 0040	24	92	-5.58	86	-0.74 67 88	-0.58 42 88	-1.14 68 89	0.08 65 103	0.36 76 108	4.13 70 110	-1.78 29 92	-1.06 29 84	98 2
AC 25 0043 109703884	2.08	RA 21 0826	AC 22 0114	25	91	-6.63	84	-0.24 62 93	-0.14 37 97	-0.24 63 95	-0.05 61 95	0.82 73 115	6.60 66 116	-1.77 21 92	-0.73 21 88	105 2
AC 25 0044 109703892	2.19	RA 21 0826	AC 21 0028	15	100	0.19	96	0.61 67 101	-0.15 40 97	0.66 67 101	-0.12 65 90	-0.86 76 89	2.47 69 105	-0.48 26 97	-0.60 25 90	99 1
AC 25 0046 109703900	2.19	RA 21 0826	AC 21 0028	14	100	0.19	96	1.12 67 106	0.00 40 100	2.26 67 111	-0.15 65 89	-0.23 76 99	4.14 69 110	-0.48 26 97	-0.60 25 90	99 1
AC 25 0047 109703918	3.78	RA 22 0028	RA 22 0121	4	107	4.61	103	0.94 67 104	0.15 40 103	2.13 68 110	0.10 65 104	1.27 76 122	4.10 69 109	4.12 29 114	1.09 29 110	93 1
AC 25 0049 109703926	1.39	B 21 0072	AC 21 0024	12	101	0.78	97	0.25 67 97	0.21 38 104	0.28 68 98	-0.09 65 93	-0.11 76 101	0.60 70 100	2.42 27 107	0.32 27 101	101 2
AC 25 0051 109703934	2.08	XT 22 0232	AC 22 0105	18	96	-2.62	91	0.24 62 97	0.12 36 103	0.26 62 98	-0.28 59 81	-0.45 72 95	-1.79 64 94	-0.49 20 96	-0.11 19 95	107 2
AC 25 0052 109703942	2.08	XT 22 0232	AC 22 0105	23	93	-4.65	88	-0.32 62 92	0.06 36 101	-0.40 62 94	-0.20 59 85	0.15 72 105	-2.67 64 92	-0.49 20 96	-0.11 19 95	107 2
AC 25 0055 109703959	2.08	B 21 0072	AC 22 0113	21	95	-3.59	89	-0.65 64 89	-0.27 36 94	-1.29 62 88	-0.04 60 95	0.87 72 116	6.56 65 116	3.25 16 111	0.59 16 104	106 2
AC 25 0058 109703983	2.08	RA 21 0826	AC 22 0107	26	89	-7.44	83	-1.10 62 85	-0.45 37 90	-1.82 63 85	-0.03 60 96	0.42 72 109	3.95 65 109	-1.77 21 92	-0.73 21 88	102 2
AC 25 0061 109704007	2.08	B 21 0072	AC 22 0101	10	102	1.10	97	0.25 61 97	0.08 32 102	0.02 62 97	-0.22 60 84	-0.64 72 92	2.19 65 104	3.25 16 111	0.59 16 104	103 2
AC 25 0065 109704023	2.08	RA 22 0028	AC 22 0108	11	102	1.01	97	-0.05 60 95	0.06 33 101	-0.16 61 95	-0.10 58 92	-0.29 71 98	-4.37 64 87	0.87 20 102	0.45 20 102	99 1
AC 25 0066 109704031	2.95	RA 22 0028	AC 21 0064	16	99	-0.47	95	0.01 67 95	0.04 41 101	0.17 67 98	0.21 64 111	1.58 76 127	2.97 69 107	1.67 30 105	0.42 29 102	106 2
AC 25 0067 109704049	2.95	RA 22 0028	AC 21 0064	7	105	3.26	101	0.92 67 104	-0.01 41 100	1.45 67 106	0.20 64 110	1.09 76 119	0.48 69 100	1.67 30 105	0.42 29 102	106 2
AC 25 0072 109704056	2.08	RA 22 0028	AC 22 0109	9	103	2.19	99	-0.29 61 92	-0.24 36 95	0.17 61 98	0.11 58 105	0.42 71 109	4.74 64 111	0.87 20 102	0.45 20 102	101 2

Young Rams



Genotyped

EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding value
Breeding values with an accuracy of below 5% are not printed

Young rams, born since 2020 with a weaning weight, and wool test, but no progeny, both parents known.

Animal Information		Parents		Rank	Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility			
Animal ID Regnr	% In- breeding	Sire ID	Dam ID		SIP	GM		Weaning Dir	Weight Mat	Body Weight	CFW	FD	SL	NLW	TWW	EPI	DAM Lambcd
						EBV	Index		EBV	Acc	Index	Acc	EBV	Acc	Index	Acc	EBV
		Active Breed Average :			101	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27		
AC 25 0077 109704072	1.64	B 21 0072	RA 22 0226	1	109	6.48	106	0.99 69 104	-0.15 41 97	1.69 69 107	0.10 67 104	-0.05 76 102	6.11 71 115	2.08 27 106	0.52 26 103	113	2
AC 25 0079 109704098	1.21	RA 22 0028	XT 18 1846	13	101	0.47	96	0.95 68 104	0.41 42 108	2.33 68 111	-0.13 65 90	0.64 76 112	-3.24 70 90	1.70 28 105	0.21 27 99	108	5
AC 25 0082 109704106	1.66	XT 22 2277	RA 22 0210	5	106	3.92	102	2.56 68 119	0.61 38 113	3.29 67 118	-0.08 65 93	0.08 76 104	0.31 69 100	-2.21 24 90	-0.03 24 96	123	2
Young rams (with known parents) average :					100	-0.10	95	0.15	0.03	0.45	-0.06	0.21	2.42	1.46	0.21		

Ewes with Progeny



Genotyped

EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding values
Breeding values with an accuracy of less than 5% are not printed

Ewes with progeny and measured at weaning, with both parents known.

Animal Information		Parents			Progeny		Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility				Rank
Animal ID Regnr	% In- breeding	Sire ID Dam ID	DAM : EPI Times lambd	Generation Progeny Weaned	1st	2nd	SIP	GM		Weaning Dir EBV Acc	Weight Mat EBV Acc	Body Weight EBV Acc	SVG EBV Acc	VD EBV Acc	SL EBV Acc	Lammers gespeen EBV Acc	TWW EBV Acc	EPI Phenotype	Times lambd	
Active Breed Average :							100.7	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27			
AC 18 0056 84461383	0.73	AC 12 0030 EN 13 0341	109 4	3	1		109	-6.73	84	-2.46 74 72	-0.74 65 84	-3.08 72 77	-0.22 70 85	-0.43 79 96	4.07 74 109	2.98 46 110	0.01 46 97	102	4	6
AC 19 0024 90203027	1.73	EN 12 0250 AC 17 0019	99 2	5			103	-8.26	82	-0.57 75 90	0.85 67 118	-1.78 71 85	-0.31 70 79	0.24 79 106	3.90 73 109	1.40 49 104	0.07 49 98	143	4	7
AC 19 0034 90202888	0.92	EN 16 0506 AC 13 0033	136 3	3			102	-9.96	79	-1.96 71 77	0.36 61 108	-2.22 68 82	-0.43 66 71	0.26 76 106	-5.80 70 84	1.55 42 104	0.35 42 101	98	3	8
AC 20 0003 92668532	1.56	EN 16 0433 AC 15 0106	125 4	3			101	-5.50	86	-0.43 69 91	-1.00 60 79	-1.59 66 86	-0.03 64 96	0.45 76 109	0.14 69 99	-2.23 38 90	-0.25 38 94	115	3	9
AC 20 0018 92668581	3.58	EN 16 0433 AC 17 0011	92 1	4			111	3.10	100	0.21 71 97	-0.64 61 86	-0.28 69 95	0.09 64 104	-0.21 76 99	4.78 70 111	1.89 40 105	0.34 40 101	128	3	3
AC 20 0021 92668607	2.89	RA 16 0738 AC 18 0031	114 3	2			93	-8.68	81	-2.11 70 75	-0.25 57 95	-2.99 66 77	-0.14 64 89	0.86 75 115	0.72 69 101	4.46 41 115	0.40 40 102	98	2	12
AC 20 0026 92668623	0.65	VM 17 0072 AC 17 0007	94 2	2			105	0.27	96	-0.50 70 90	0.76 57 116	-0.56 67 93	-0.17 64 88	0.13 75 104	-5.53 69 84	2.55 42 108	1.16 42 111	103	2	6
AC 21 0010 94536547	0.41	VM 18 0551 AC 16 0106	119 5	2			107	-5.97	86	-1.52 69 81	-0.28 57 94	-2.99 66 77	0.21 64 111	0.10 75 104	-2.80 69 91	-1.38 41 93	-1.14 41 83	101	2	3
AC 21 0024 94536760	1.03	VM 18 0551 AC 17 0036	119 4	1			103	-4.41	88	-0.79 69 88	0.37 53 108	-1.06 68 90	-0.01 66 97	0.38 77 108	-3.61 71 89	-1.21 43 94	-0.34 42 93	101	2	8
AC 21 0025 94536786	0.50	WW 18 047 AC 18 0009	111 3	3			102	-8.30	82	-1.33 71 82	-0.03 59 99	-2.49 69 81	-0.08 67 93	0.40 78 108	1.44 72 103	0.22 39 99	-0.46 39 91	105	2	9
AC 21 0028 94536828	1.89	RA 18 0346 AC 19 0106	99 2	2			97	-2.74	91	0.56 65 100	0.86 52 118	1.88 65 108	-0.45 62 70	0.15 75 104	-0.66 68 97	3.03 28 110	0.47 28 102	99	1	18
AC 21 0037 94536927	2.29	RA 18 0346 AC 18 0037	101 3	2			96	-9.67	79	-1.53 70 81	-0.40 56 92	-2.24 68 82	-0.16 65 88	0.39 76 108	-0.18 70 98	-1.58 37 92	-0.61 37 89	99	2	20
AC 21 0039 94536943	2.67	RA 18 0346 AC 19 0017	92 1	1			102	-3.31	90	0.89 67 103	0.30 49 106	1.76 66 108	-0.31 63 79	0.67 75 113	-2.12 68 93	2.49 33 108	0.26 33 100	93	1	10
AC 21 0040 94536968	0.63	WW 18 047 AC 18 0031	114 3	2			98	-8.73	81	-1.64 70 80	-0.35 56 93	-2.12 68 83	0.05 65 101	1.27 77 122	-0.03 70 99	0.43 39 100	-0.44 39 91	98	2	17
AC 21 0041 94536984	1.91	VM 18 0551 AC 18 0056	102 4	2			110	-1.82	92	-1.62 70 80	-0.69 57 85	-1.95 69 84	0.16 66 108	0.17 77 105	6.68 71 116	2.62 40 108	-0.20 40 94	96	1	2
AC 21 0043 94537008	0.56	WW 18 047 AC 17 0007	94 2	3			93	-9.94	79	-0.56 71 90	0.15 59 103	-1.26 69 88	-0.16 67 88	1.75 78 129	2.80 72 106	3.08 38 110	0.19 38 99	105	2	22

Ewes with Progeny



Genotyped

EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding values
Breeding values with an accuracy of less than 5% are not printed

Ewes with progeny and measured at weaning, with both parents known.

Animal Information		Parents			Progeny		Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility				Rank	
Animal ID Regnr	% In- breeding	Sire ID Dam ID	DAM :		Generation Progeny Weaned 1st 2nd	SIP	GM	Weaning Weight		Body Weight	SVG		VD		SL	Lammers gespeen	TWW		EPI	Times lambd	
			EPI	Times lambd				Dir	Mat		Acc	Index	Acc	Index			Acc	Index			
		Active Breed Average :				100.7	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27					
AC 21 0045 94537040	3.28	VM 18 0551 AC 15 0014	130	5	2	99	-3.85	89	-1.10 72	0.95 85	-1.61 70	0.04 68	1.10 78	1.11 72	4.21 41	0.56 41	97	1	14		
AC 21 0046 94537065	2.67	RA 18 0346 AC 19 0016	96	2	3	99	-3.92	89	0.68 71	0.89 59	1.51 68	-0.50 65	0.27 76	-7.14 70	3.53 37	0.62 37	109	2	13		
AC 21 0047 94537081	3.45	RA 18 0346 AC 16 0027	109	4	2	97	-13.46	73	-1.43 71	-0.35 57	-1.95 65	-0.34 63	0.37 74	-1.08 68	-1.62 39	-1.04 39	96	2	19		
AC 21 0062 94537180	0.45	WW 18 047 AC 16 0110	111	4	1	93	-12.26	75	-0.79 67	-0.29 51	-1.95 64	-0.23 62	0.82 74	-4.49 67	0.15 35	-0.72 34	93	1	21		
AC 21 0064 94537222	1.13	VM 18 0551 AC 17 0032	125	4	2	101	-3.39	90	-0.58 71	-0.05 58	-1.23 70	0.21 68	1.52 78	0.77 73	2.07 43	0.16 43	106	2	11		
RA 22 0116 94738085	2.32	RA 20 0051 RA 20 0604	89	1	2	99	2.91	100	-0.06 70	-0.36 55	0.74 70	-0.04 68	0.22 78	-2.02 72	5.83 38	0.84 37	97	1	144		
RA 22 0121 94738184	3.67	RA 20 0051 RA 20 1044	123	4	1	95	3.41	101	0.34 71	0.10 56	1.66 72	0.11 70	1.85 79	3.84 74	6.95 41	1.49 41	93	1	205		
RA 22 0143 94738382	1.80	RA 20 0051 EN 14 0059	141	7	1	99	1.68	98	0.31 73	-0.25 58	0.56 72	0.00 70	0.69 78	1.78 73	5.31 45	0.88 45	91	1	149		
RA 22 0210 94745544	4.56	RA 20 0325 EN 16 0638	125	5	2	96	0.61	96	1.61 72	0.80 60	4.16 71	-0.02 69	1.36 78	8.63 73	-6.93 43	-0.35 43	123	2	191		
RA 22 0223 94738788	3.23	RA 20 0325 RA 15 1149	127	6	1	96	-0.60	94	1.03 71	0.62 56	2.33 71	-0.05 69	0.90 79	8.17 73	-3.96 40	-0.16 40	95	1	199		
RA 22 0226 94738846	2.52	RA 20 0325 RA 20 0506	122	4	2	94	0.21	96	0.68 72	0.30 59	2.15 72	-0.02 70	0.94 79	9.94 74	-1.88 42	0.06 42	113	2	213		
RA 22 0410 94739687	1.89	RA 20 0621 RA 19 1307	136	5	1	95	-7.92	82	-1.25 68	0.21 50	-0.81 67	-0.49 65	-0.53 76	-8.31 70	-0.61 26	-0.27 25	92	1	203		
RA 22 0923 94742707	7.89	RA 20 0861 RA 20 1032	116	4	2	90	-7.66	83	-1.58 71	-1.04 55	-2.19 70	-0.10 68	0.67 78	3.74 72	3.39 36	-0.05 36	93	1	242		
RA 22 1033 94743366	2.83	LM 20 0195 RA 19 0455	120	5	1	94	-3.53	90	0.24 70	-0.22 53	-0.17 69	-0.12 66	0.51 77	1.16 71	-1.14 36	0.11 36	92	1	215		
RA 22 1038 94743408	1.13	LM 20 0195 RA 19 0933	95	3	1	96	-2.19	92	-0.24 70	-0.62 53	0.61 69	-0.15 67	0.21 77	-3.24 71	1.03 37	-0.06 37	90	1	192		
RA 22 1048 94743481	0.85	LM 20 0195 RA 17 0713	101	4	2	96	-3.64	89	-0.12 72	0.27 58	0.26 70	-0.43 67	-0.60 77	-7.24 71	1.42 41	0.10 41	114	2	185		

Ewes with Progeny



Genotyped

EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding values
Breeding values with an accuracy of less than 5% are not printed

Ewes with progeny and measured at weaning, with both parents known.

Animal Information		Parents			Progeny		Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility				Rank
Animal ID Regnr	% In- breeding	Sire ID Dam ID	DAM :		Generation Progeny Weaned 1st 2nd	SIP	GM	Weaning Dir	Weight		Body Weight	SVG	VD	SL	Lammers gespeen	TWW	EPI	Times lambd		
			EPI	Times lambd					Acc	Index										Acc
		Active Breed Average :				100.7	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27				
RA 22 1103 94743564	2.00	RA 20 0234 RA 20 1404	92	2	1	90	-7.21	84	-0.54 65	0.19 49	-0.38 64	-0.36 61	0.47 73	-4.57 66	-0.20 29	0.16 29	90	1	245	
RA 22 1105 94743580	2.00	RA 20 0234 RA 20 1433	89	1	1	92	-5.49	86	-0.42 63	0.44 47	-0.28 63	-0.31 60	0.35 72	-1.75 65	-0.24 29	0.25 29	92	1	234	
RA 22 1107 94743606	2.00	RA 20 0234 RA 20 1429	123	4	1	90	-6.98	84	-1.23 68	-0.07 52	-1.48 66	-0.23 63	0.53 75	3.03 68	-0.24 29	0.25 29	91	1	244	
RA 22 1113 94743663	2.00	RA 20 0234 RA 20 1408	118	4	2	92	-4.09	89	-0.48 68	0.78 55	0.53 67	-0.14 64	0.88 75	1.90 69	-1.61 33	0.06 33	110	2	230	
RA 22 1119 94743721	2.00	RA 20 0234 RA 20 1402	112	4	2	94	-5.29	87	-0.63 68	0.53 55	-1.12 68	-0.40 63	-0.43 75	-4.59 68	1.55 28	0.34 28	98	1	218	
XT 18 1846 82859216	1.47	XT 15 6532 XT 14 1464	104	2	4 1	109	-5.16	87	-0.01 71	0.81 60	2.69 71	-0.37 68	0.81 79	-6.56 73	2.11 37	-0.27 37	108	5	8	
XT 20 5016 91040733	1.44	XT 18 0808 XT 18 5819	119	6	2 1	106	-5.54	86	-1.24 72	0.43 56	-1.95 71	-0.14 69	-0.45 79	0.65 74	-1.57 40	-0.52 39	97	3	18	
XT 21 1119 93126514	2.71	XT 18 1804 XT 16 1628	109	4	2	99	-3.95	89	-0.67 71	0.30 57	-0.86 70	-0.15 67	0.31 78	-0.89 72	3.44 37	0.21 37	94	2	75	
XT 21 1152 93129153	1.59	XT 19 4920 XT 19 1925	89	3	1	101	-10.24	79	-0.15 69	-0.74 50	-0.52 70	-0.22 68	0.44 78	-3.38 72	-2.11 39	-1.27 39	86	3	54	
XT 21 2145 93125698	1.47	XT 19 5948 XT 14 2467	169	5	1	107	-6.17	85	-2.35 67	-1.20 50	-4.51 65	0.05 62	-0.38 74	0.81 67	1.53 31	-0.27 31	86	2	15	
XT 22 0226 95809513	1.75	XT 20 5023 XT 17 0773	96	3	2	106	0.87	97	-0.05 66	0.45 52	-0.76 67	-0.15 64	-0.46 76	6.27 70	2.64 23	0.74 23	102	1	17	
XT 22 5247 95808853	1.67	XT 19 0966 XT 15 5573	188	7	1	96	-6.67	84	-0.19 68	-0.43 50	-0.99 66	0.13 64	1.94 75	1.78 69	1.51 26	-0.07 25	95	1	100	
Ewes with progeny average :						98.9	-4.8	87	-0.58	0.04	-0.65	-0.15	0.48	0.05	1.02	0.04				

Young Ewes



EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding values
Breeding values with an accuracy of less than 5% are not

Ewes born since 2020 with a weaning weight and wool test, no progeny, with both parents known.

Animal Information		Parents			Selection Indices			EBV - Weights						EBV - Wol						EBV - Fertility				Rank
Animal ID Regnr	% In- breeding	Sire ID Dam ID	DAM REPROD :		SIP	GM	Weaning Weight				Body Weight		CFW		FD		SL	NLW		TWW				
			EPI	Times lambded			Dir	Mat	Weight	Acc	Index	Acc	Index	Acc	Index	Acc	Index	Acc	Index	Acc		Index	Acc	Index
							EBV	Index		EBV	Index	EBV	Index	EBV	Index	EBV	Index	EBV	Index	EBV	Index			
		Active Breed Average :			100.7	2.8	100			0.54	0.01	0.56		0.03	-0.14	0.50		0.46	0.27					
AC 25 0004 109703322	1.61	RA 22 0028 XT 22 0226	102	1	106	4.36	103			1.55 67	0.18 39	1.70 107		0.07 64	0.61 75	6.03 115		1.96 23	0.71 105		2			
AC 25 0010 109703330	1.57	B 21 0072 XT 22 5247	95	1	95	-3.92	89			0.14 68	0.00 37	-0.05 100		-0.10 66	1.13 76	4.62 111		3.78 21	0.45 102		23			
AC 25 0012 109703355	2.21	B 21 0072 XT 21 1182	99	1	99	-0.48	95			-0.57 63	-0.15 32	-1.04 97		-0.14 61	-0.31 72	2.20 104		3.25 16	0.59 104		17			
AC 25 0014 109703363	2.21	B 21 0072 XT 21 4178	100	1	97	-2.44	91			-0.28 63	0.00 32	-0.44 100		-0.13 61	0.55 72	-0.33 98		3.25 16	0.59 104		22			
AC 25 0015 109703371	2.21	B 21 0072 XT 21 0168	109	1	101	0.37	96			0.51 64	0.30 35	1.11 106		-0.12 62	0.52 73	0.22 99		3.25 16	0.59 104		13			
AC 25 0022 109703397	2.21	XT 22 2277 XT 21 1192	101	1	98	-1.27	93			-0.12 62	0.33 28	-0.31 107		-0.17 59	-0.26 72	4.39 110		1.48 10	0.25 100		19			
AC 25 0026 109703439	2.08	B 21 0072 AC 22 0111	100	1	103	2.05	99			-0.12 63	0.06 34	-0.49 101		-0.14 60	-0.78 72	-4.02 88		3.25 16	0.59 104		7			
AC 25 0029 109703454	1.54	XT 22 5209 AC 21 0045	97	1	103	2.14	99			-0.68 67	0.20 37	-1.63 104		0.23 64	0.29 75	3.71 108		2.73 23	0.60 104		6			
AC 25 0030 109703462	1.54	XT 22 5209 AC 21 0045	97	1	100	-0.09	95			-1.03 67	0.34 37	-2.15 107		0.11 64	0.17 75	-0.21 98		2.73 23	0.60 104		14			
AC 25 0033 109703488	2.08	XT 22 5209 AC 22 0100	101	1	100	-0.46	95			-0.85 60	-0.03 28	-2.08 99		-0.02 58	-0.44 71	-6.71 81		0.85 11	0.42 102		16			
AC 25 0034 109703496	1.33	RA 22 0028 AC 21 0043	105	2	98	-1.55	93			0.21 67	0.25 41	0.90 105		-0.09 64	0.95 75	-0.66 97		2.18 28	0.44 102		20			
AC 25 0038 109703504	2.71	RA 21 0826 RA 22 0223	95	1	104	2.53	100			1.64 68	-0.36 42	2.21 92		0.17 67	0.16 76	7.03 117		-3.97 29	-0.92 86		5			
AC 25 0045 109703520	1.45	B 21 0072 AC 21 0037	99	2	101	0.57	96			0.20 68	-0.23 39	-0.03 95		-0.18 65	-0.73 76	1.28 102		2.23 25	0.19 99		10			
AC 25 0048 109703538	1.89	B 21 0072 AC 19 0103	119	4	104	3.00	100			-0.09 65	-0.29 38	-1.03 94		-0.05 60	-0.91 72	-5.49 84		3.25 16	0.59 104		3			
AC 25 0050 109703546	2.08	B 21 0072 AC 22 0104	101	2	101	0.48	96			-0.85 63	-0.12 35	-1.36 97		-0.29 60	-1.40 72	-3.96 88		3.25 16	0.59 104		12			
AC 25 0053 109703553	1.37	RA 22 0028 AC 21 0025	105	2	103	1.97	99			0.54 67	-0.11 41	0.50 98		0.08 64	0.19 75	0.34 100		0.75 28	0.11 98		8			

Young Ewes



EBV - Estimated Breeding Value

*Inbreeding in red is high

Indices in top 10%
Above breed average
Indices in bottom 10%

Acc: Accuracy of the breeding values
Breeding values with an accuracy of less than 5% are not

Ewes born since 2020 with a weaning weight and wool test, no progeny, with both parents known.

Animal Information		Parents			Selection Indices			EBV - Weights			EBV - Wool			EBV - Fertility		Rank
Animal ID Regnr	% In- breeding	Sire ID Dam ID	DAM REPROD :		SIP	GM		Weaning Weight		Body Weight	CFW	FD	SL	NLW	TWW	
			EPI	Times lambd				Dir	Mat							
								EBV	EBV	EBV	EBV	EBV	EBV	EBV	EBV	
								Acc	Index	Acc	Index	Acc	Index	Acc	Index	
Active Breed Average :					100.7	2.8	100	0.54	0.01	0.56	0.03	-0.14	0.50	0.46	0.27	
AC 25 0054 109703561	1.37	RA 22 0028 AC 21 0025	105	2	92	-5.46	86	-0.88 67	0.34 41	-0.28 67	-0.07 64	1.26 75	3.16 69	0.75 28	0.11 28	27
AC 25 0060 109703579	1.93	RA 21 0826 RA 22 1113	110	2	99	-1.09	94	0.49 68	-0.15 42	1.02 68	0.07 66	0.32 76	5.08 70	-2.79 27	-0.81 27	18
AC 25 0062 109703587	2.91	RA 21 0826 AC 20 0003	115	3	94	-4.22	88	-0.18 68	-1.01 43	-1.15 68	0.11 65	0.20 76	3.74 70	-3.11 28	-0.96 28	25
AC 25 0063 109703595	5.08	RA 22 0028 RA 22 1119	98	1	101	0.68	97	0.42 67	0.28 40	0.37 67	-0.12 64	-0.01 75	0.18 69	1.42 25	0.51 25	9
AC 25 0068 109703603	2.08	B 21 0072 AC 22 0110	102	2	104	2.83	100	0.08 63	-0.40 35	-0.54 63	0.05 60	-0.18 72	5.43 65	3.25 16	0.59 16	4
AC 25 0070 109703629	2.58	RA 22 0028 AC 20 0018	128	3	111	7.57	108	1.54 67	-0.46 42	1.86 67	0.23 64	0.31 75	4.74 69	1.59 29	0.51 28	1
AC 25 0074 109703645	2.08	B 21 0072 XT 22 0062	102	1	100	-0.27	95	0.40 63	0.10 32	0.30 63	-0.10 61	0.41 72	-0.10 66	3.25 16	0.59 16	15
AC 25 0075 109703652	1.48	RA 22 0028 XT 20 5016	97	3	94	-4.07	89	-0.29 68	0.48 40	0.16 68	-0.07 65	1.01 76	1.88 69	-0.14 29	0.08 28	24
AC 25 0076 109703660	1.86	B 21 0072 RA 22 0410	92	1	94	-4.59	88	-0.05 68	0.35 37	0.06 68	-0.50 66	-0.50 76	-6.13 70	2.72 21	0.35 20	26
AC 25 0080 109703678	1.21	RA 22 0028 XT 18 1846	108	5	98	-1.74	93	0.75 68	0.67 42	2.67 68	-0.21 65	1.04 76	4.00 70	1.70 28	0.21 27	21
XT 22 7227 95811329	1.52	XT 20 5023 XT 18 7821	122	5	112	2.34	99	0.49 68	0.25 44	-0.21 68	-0.14 66	-0.77 76	4.17 71	1.69 28	0.54 27	3
Young Ewes (with known parents) average :					100.4	0.0	95	0.11	0.03	0.00	-0.06	0.11	1.28	1.65	0.30	

