

Boerboel Genetic Evaluation

SA Stud Book
February 2026

0 0 JENA BOERBOELS

4044 MAINWAY MEADOWS W/F

0

HARARE

0002



Kennel Number: 719361



Estimated Breeding Values (EBVs)

Litter size and conformation traits are influenced by many genes, with every gene making a small contribution to the expression of the trait. These traits are, however, also influenced by physiological and physical environmental factors, for example the age and sex of the animal, the feeding and management the animal received and year x season effects. It is therefore not possible to know what the animal's genetic potential for the trait is by merely using the measurement.

With BLUP (*Best Linear Unbiased Prediction*) methodology an animal's genetic potential is predicted by making use of the genetic parameters (heritabilities and genetic correlations) of the breed, pedigree information, as well as the measurements of the animal as deviation from the average of the animals that were measured with the animal. Adjustments are made for differences in sex, age, classifier, management and year x season effects to ensure that fair comparisons are made amongst the animals. These predictions are called **EBVs (Estimated Breeding Values)** which indicates the animal's genetic potential for the specific trait and predicts the type of progeny that the animal will breed.

Pedigree information determines the genetic level at which the animal enters the genetic evaluation, it links the information of all relatives together and it is used to determine the genetic ranking of the breed. For young animals that were not yet measured, mid-parent EBVs are estimated, which are the average of the EBVs of the sire and dam. These breeding values are still inaccurate and can change a lot when more information (measurements) is included in subsequent genetic evaluations.

Breeding values therefore enables a breeder to compare his/her animals genetically to one another, as well as to all other animals of the breed. This Report gives a summary of the genetic levels of the active males and females of the relevant breeder in comparison to that of the breed for linearly classified traits and litter size. The genetic change that occurred over time can also easily be seen.

Breeding Value Indices (EBVi)

For easy interpretation, EBVs are standardised and expressed relative to the average (100) of the active animals of the breed, using 12 as standard deviation (SD). This means that an EBV of 100 is equal to the active average of the breed, and more or less 50% of the animals will have EBVi's of between 90 and 110, which indicates an average animal. Only 0.5% of the EBVi's will be lower than 64 and 0.5% will be higher than 136. Higher EBVi's indicate better genetic potential for the trait.

Accuracy

EBVs are published with an **accuracy value (%)**, which indicates how much info was available for the estimation of the EBV and with how much trust it can therefore be applied. Factors that influence the accuracy include heritability, completeness and correctness of pedigree information, number of own and measured relatives and the size of the group in which the animal was measured. When an accuracy of 85% and more is reached, the EBV should not change significantly with the inclusion of more information in subsequent genetic evaluations.

Kennel and Breed Averages: The different traits' averages of the breeder's active animals are compared to those of the breed in the Kennel and Breed Average Graphs and Tables. The breeder can therefore easily see for which traits the animals are above or below the average of the breed.


Genetic Trends: The Genetic Trends indicate the genetic change that occurred over time in the animals bred by the breeder compared to that of the breed. These are calculated as the average of the breeding values of all measured animals bred by the breeder, per year of birth.

Inbreeding: Inbreeding is the mating of related animals and is widely viewed as being harmful. The danger of inbreeding lies there in that it can gradually decrease fertility and survivability – a phenomenon known as inbreeding depression. It can cause lower conception rates, more abortions, more stillborn and weak offspring and a higher susceptibility to diseases. Indications in dogs include a low litter size as well as problems relative to fertility, such as invisible heat, behavioural problems when mating and low sperm quality. Overall, reduced litter size most likely reflects inbreeding depression in several traits, both physiological and behavioural, as well as a high frequency of non-viable embryos due to lethal alleles. Generally, the effects of inbreeding can become noticeable at a value of 6.25%, therefore it is generally recommended that individuals should have inbreeding coefficients of less than 6.25%. Breeders should therefore avoid matings causing inbreeding of more than 6.25% in the progeny. The theoretical maximum inbreeding coefficient is 50%. However, it should be noted that not all inbred animals show signs of inbreeding depression.

The data used in this evaluation was obtained from LOGIX from linear classifications that were done by the Breeder Society. All possible attempts have been made to present the information correctly. SA Stud Book does not accept responsibility for the use and interpretation of the information presented on animals in this publication.

Genetic Levels

Kennel Average relative to Breed Average, where Breed Average is indicated as 0



Average Breeding Value Indices (EBVi) where Breed Average = 100

Trait	All Animals	Males	Females
Litter size (LS)	87	85	88
Movement (Mv)	102	98	103
Temperament (Tmp)	108	106	109
Coat (Ct)	103	109	101
General Appearance	116	120	114
Head & Face (H&F)	109	110	108
Front Quarter (FQ)	110	110	110
Centre (C)	113	113	113
Hind Quarter (HQ)	111	109	113
Final Score (FS)	112	112	112
Hip Health (HH)	107	106	108
Elbow Health (ELM)	102	101	102

Inbreeding

Notes

Inbreeding primarily leads to decreased fertility, low survivability of pups and low disease resistance.

Inbreeding coefficients are calculated for all animals of the breed and indicates the probability that two genes of a pair are the same due to a common ancestor in the pedigree of the animal.

The adverse effects of inbreeding can be observed from 6.25% onwards.


It is therefore highly recommended to record pedigree information as accurately. Inbreeding is dependent on pedigree information. Conventional methods for the calculation of inbreeding coefficients usually assume that unknown parents are unrelated, which is usually far from the truth. Inbreeding coefficients are therefore conservative indicators of the level of inbreeding.

Incomplete or incorrect pedigree information can be reported to SA Stud Book via the Breed Society.

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Rate of Inbreeding




Inbreeding Levels


	All Animals		Males		Females		Sires		Dams	
	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed
Number	7	49007	2	25141	5	23857	4	3371	4	6287
Average Inbreeding	8.67	8.49	12.42	8.46	7.17	8.53	7.63	8.01	5.41	8.00
Number inbred	7	48727	2	24990	5	23728	4	24990	4	23728
Number >6.25% Inbr.	7	30600	2	15537	5	15058	2	15537	2	15058

Genetic Trends


Litter Size




Movement




Temperament




Coat



General Appearance




Head & Face




Genetic Trends


Front Quarter



Centre




Hind Quarter




Final Score & Health Traits


Final Score



Hip Health



Elbow Health



Estimated Breeding Value Indices (EBVi's)

Males

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	Sire	Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
JENA CHUBBS	KING GRANITE SANITIZER		17.56	106	104	77	111	111	109	125	108	114	119	113	116	7	1				
	2023	BANTU MAKENZI JENA		5	5		14	5	44	12	43	16	22	27	36						
JENA G SIX	DTK JOHN WICK		7.28	107	97	94	86	102	109	116	112	107	108	105	109						
	2025	BANTU ROXY		5	5		19	5	35	18	34	20	22	25	30						

Estimated Breeding Value Indices (EBVi's)

Females

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information

Inbreed

Health

Fertility

General

Appearance

Progeny :

Appraisal

Health

Progeny with litters

Sire		%	Hips	Elbows	Litter Size	Move-	Tempe-	Coat	General	Head &	Front	Centre	Hind	Final	Progeny Litters	Progeny	Hips	Elbows	with	
BYr	Dam		Acc.	Acc.	Acc.	ment	rament		Appear.	Face	Quarter		Quarter	Score		Kennels				
A	2021	ALESSANDRO BENJIE	6.36	104	101	89	119	114	106	120	105	113	116	116	115	5	1			
		KING GRANITE STEMBENI		5	5	21	35	5	52	34	51	35	38	41	46					
	2022	ALESSANDRO BENJIE	6.36	104	101	72	112	111	97	116	104	110	119	118	114	3	2	3		1
				KING GRANITE HAVOCKESS	5	5	27	33	5	50	33	49	34	37	39					
	2024	JAMBAL ENGINE	8.55	116	114	89	112	116	83	102	109	114	112	120	113					
				BANTU MAKENZI JENA	5	5	5	33	5	50	32	49	34	36	39					
2025	DTK JOHN WICK	7.28	107	97	94	86	102	109	116	112	107	108	105	109						
			BANTU ROXY	5	5	5	19	5	35	18	34	20	22	25						30
2025	DTK JOHN WICK	7.28	107	97	94	86	102	109	116	112	107	108	105	109						
			BANTU ROXY	5	5	5	19	5	35	18	34	20	22	25						30

Estimated Breeding Value Indices (EBVi's)

Sires

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Sire Information		Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels	Hips	Elbows	
ALESSANDRO BENJIE		CDH MONTARO	6.16	100	95	93	113	116	107	125	110	118	125	123	121	44 11	15 12	1	1	10 4
	2019	KLEIN SANDFONTEIN ZUZU		5	5	32	50	24	64	49	63	50	53	55	59					
KING GRANITE SANITIZER		IVANOK KING	4.45	107	107	81	120	112	106	123	102	111	113	110	111	21 7	5 5			3 2
	2020	GIANTS OF AFRICA HAVOCKESS		5	5	28	41	10	57	40	56	42	44	47	52					
DTK JOHN WICK		ELEVATION TRIOMF	6.71	111	94	99	52	91	113	112	118	102	99	93	103	26 9	7 6			1 1
	2021	MARVILLOSO GRIZELDA BLANCA		5	5	7	42	12	58	41	57	43	45	47	52					
JAMBAL ENGINE		MIDDELPOS BBD GAMKA	13.19	128	128	106	102	119	81	103	123	124	112	117	118	8 3	3 2			
	2021	JAMBAL DOUBLE TROUBLE		5	5	29	55	31	67	54	66	55	57	59	63					

Estimated Breeding Value Indices (EBVi's)

Dams

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe- r ament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
KING GRANITE STEMBENI	2017	IVANOK HEROD	4.45	107	107	82	120	109	97	105	103	106	108	114	109	10	5	4		2	
		GIANTS OF AFRICA HAVOCKESS		5	5	39	34	5	49	34	48	35	37	40	44			3		1	
KING GRANITE HAVOCKESS	2020	IVANOK KING	4.45	107	107	65	117	109	96	116	97	104	112	110	107	8	3	3		2	
		GIANTS OF AFRICA HAVOCKESS		5	5	39	44	13	62	43	60	45	47	50	55			3		1	
BANTU ROXY	2021	ALESSANDRO BENJIE	6.36	104	101	89	119	114	106	120	105	113	116	116	115	5	1				
		KING GRANITE STEMBENI		5	5	21	35	5	52	34	51	35	38	41	46						
BANTU MAKENZI JENA	2022	ALESSANDRO BENJIE	6.36	104	101	72	112	111	97	116	104	110	119	118	114	3	2	3		1	
		KING GRANITE HAVOCKESS		5	5	27	33	5	50	33	49	34	37	39	44			2		1	

