

logix



SA STAMBOEK
STUD BOOK

Logix *Melk/Milk*

Kudde Gesondheids Verslag / Herd Health Report

HYDE

POSTNET SUITE 508
PRIVATE BAG H607
HEIDELBERG
1438

firmhyde@gmail.com

Test-day: 2024-12-11

Jersey Cows

LogixMilk Technical Advisor:

SURETHA FRANCIS

083 584 0814

suretha@studbook.co.za

December 21, 2024

List of Tables

1	Summary for Test-day 2024-12-11 (Number of Samples Tested: 12)	2
2	Quantification of Specific Cases (% of herd) and Loss in Production on 2024-12-11	2
3	Benchmark: Herd with the Top 25%, Average and Bottom 25% for different traits tested the last month of for Jersey Cows	2
4	Summary of Somatic Cell Counts per Lactation Group in the herd for 12 Jersey Cows	3
5	Somatic Cell Count (SCC) Analysis: Percentage (Number) of Cows within Stage of Lactation and Lactation Group for SCC > 200k and SCC > 750k for 12 Jersey Cows	3
6	Bulk Tank Results for 2024-12-11 (Number of Samples Tested: 0)	3
7	Somatic Cell Count (SSC) distribution and Milk Loss Monitor 2024-12-11	5
8	Cows Under Pressure (SCC > 500k and Lactose % < 4.6)	6
9	Cows with Somatic Cell Counts > 200k (Status: N=New -, C=Chronic -, R=Recurring - and CR=Chronic and Recurring Infections)	7
10	Possible Ketosis Cases; Cows under 100 days in milk (Prot. Fat Ratio < 0.68 for Jerseys and < 0.74 for other Breeds)	8
11	Possible Acidosis Cases (Protein % > Butterfat %)	9
12	Dry Cows with a Dry Period more than 55 Days	10
13	Cows with an active Inter-Calving Period (ICP) > 400 days	11

List of Figures

1	Somatic Cell Count (SCC) Trend	3
2	Percentage (%) of Cows with Somatic Cell Counts (SCC) more than 200k and 750k, respectively, per Test-day	4
3	Somatic Cell Count (SCC) Monitor Trend	4
4	Average Somatic Cell Counts (SCC, x1000/ml) per Day In Milk (DIM)	5
5	Somatic Cell Count (SCC) Distribution for Test-day (2024-12-11)	5

HERD HEALTH SUMMARY REPORT

HYDE - Jersey Cows

Table 1: Summary for Test-day 2024-12-11 (Number of Samples Tested: 12)

Analysis	Results	Norms*	% Cows outside norms*
Milk Production (kg)	Avg.: 19.7 Median: 18.2		
Butterfat (%)	Avg.: 3.93 Median: 3.98	3.5 % - 6.0 %	Too Low: 25 % Too High: 0 %
Protein (%)	Avg.: 3.62 Median: 3.64	3.0 % - 4.5 %	Too Low: 0 % Too High: 0 %
Lactose (%)	Avg.: 4.65 Median: 4.66	> 4.7 %	Too Low: 67 %
MUN (mg/dl)	Avg.: 15.61 Median: 14.65	8 - 16 mg/dl	Too Low: 0 % Too High: 42 %
SCC (x1000/ml)	Avg.: 93 Median: 41 % < 100 : 75.0 % % < 200 : 83.3 % % < 500 : 100.0 % % > 750 : 0.0 % % > 1 000 : 0.0 % % > 1 500 : 0.0 %	< 200 > 64 % > 75 % > 92 % < 4.8 % < 1 % NONE	Too High: 0 % Too High: 0 % Too High: 0 % Too High: 0 % Too High: 0 % Too High: 0.0 %
Butterfat Protein ratio	Average: 1.1	> 1.0	Too High: 33.3 %

Table 2: Quantification of Specific Cases (% of herd) and Loss in Production on 2024-12-11

Loss in Production due to:		Avg/Median SCC (Number (n)):	
SSC (kg per day):	4.7 (2.0 %)	1st Lact Cows:	39/39 (2)
DIM (kg per day):	14.2 (6.0 %)	2nd Lact Cows:	26/27 (2)
		3rd Lact Cows:	38/27 (3)
No. of Cows Under Pressure:	0 (0.0 %)	4th Lact Cows:	26/26 (1)
No. Cows with SCC > 200k:	2 (16.7 %)	5th Lact Cows:	0/ (0)
No. Cows with SCC > 750k:	0 (0.0 %)	6+ Lact Cows:	210/184 (4)
No. New Infections:	0 (0.0 %)		
No. Recurring Infections:	2 (16.7 %)	Avg. SCC for SCC > 750k (n):	(0)
Cure Rate for SCC > 200k:	0.0 % (0)		
Cure Rate for SCC > 750k:	0.0 % (0)	Avg. Days In Milk (DIM) for Herd:	186
No. Chronic Infections (SCC > 200k):	0 (0.0 %)	Median of DIM for Herd:	154
No. Chronic Infections (SCC > 750k):	0 (0.0 %)		
		Avg. DIM for SCC < 200k (n):	179 (10)
No. of Possible Ketosis Cases:	0	Avg. DIM for SCC > 200k and < 750k (n):	226 (2)
No. of Possible Acidosis Cases:	4	Avg. DIM for SCC > 750k (n):	(0)
Avg. Age at First Calf (Months):	26	Avg. Parity for Herd:	4
Avg. Inter-Calf Period (Days):	532	Avg. Parity for SCC < 200k:	4
No. Flushings:	NA	Avg. Parity for SCC > 200k and < 750k:	8
No. Embrios:	NA	Avg. Parity for SCC > 750k:	

Table 3: Benchmark: Herd with the Top 25%, Average and Bottom 25% for different traits tested the last month of for Jersey Cows

	Milk Production (kg)	Butterfat (%)	Protein (%)	Lactose (%)	SCC (x1000/ml)	MUN (mg/dl)
Herd	19.7	3.93	3.62	4.65	93	15.61
Top 25%	27.0	5.66	4.15	4.82	37	13.90
Breed Avg	22.1	5.07	3.86	4.64	289	11.01
Bottom 25%	16.6	4.55	3.60	4.53	251	8.20

*** PLEASE CONSULT YOUR VETERINARIAN, PROFESSIONAL ANIMAL SCIENTIST or LogixMilk TECHNICAL ADVISOR for the interpretation of these results and professional advisory services.**

Table 4: Summary of Somatic Cell Counts per Lactation Group in the herd for 12 Jersey Cows


Lactation Group	% of Herd	Early Lactation	Mid Lactation	Late Lactation	% SCC > 200k	% SCC > 750k
1	16.7% (2)	0.0% (0)	16.7% (2)	0.0% (0)	0.0% ()	0.0% ()
2	16.7% (2)	0.0% (0)	16.7% (2)	0.0% (0)	0.0% ()	0.0% ()
3	25.0% (3)	8.3% (1)	0.0% (0)	16.7% (2)	0.0% ()	0.0% ()
4	8.3% (1)	0.0% (0)	0.0% (0)	8.3% (1)	0.0% ()	0.0% ()
5	0.0% (0)	0.0% (0)	0.0% (0)	0.0% (0)	0.0% ()	0.0% ()
6+	33.3% (4)	0.0% (0)	16.7% (2)	16.7% (2)	16.7% (2)	0.0% ()
Herd	100% (12)	8.3% (1)	50.1% (6)	41.7% (5)	16.7% (2)	0% (0)

Table 5: Somatic Cell Count (SCC) Analysis: Percentage (Number) of Cows within Stage of Lactation and Lactation Group for SCC > 200k and SCC > 750k for 12 Jersey Cows

Lactation Group	Early Lactation		Mid Lactation		Late Lactation	
	% SCC > 200k	% SCC > 750k	% SCC > 200k	% SCC > 750k	% SCC > 200k	% SCC > 750k
1	0% ()	0% ()	0.0% ()	0.0% ()	0% ()	0% ()
2	0% ()	0% ()	0.0% ()	0.0% ()	0% ()	0% ()
3	0.0% ()	0.0% ()	0% ()	0% ()	0.0% ()	0.0% ()
4	0% ()	0% ()	0% ()	0% ()	0.0% ()	0.0% ()
5	0% ()	0% ()	0% ()	0% ()	0% ()	0% ()
6+	0% ()	0% ()	50.0% (1)	0.0% ()	50.0% (1)	0.0% ()

Table 6: Bulk Tank Results for 2024-12-11 (Number of Samples Tested: 0)

Tank No	Butterfat (%)	Protein (%)	Lactose (%)	SCC (x1000/ml)	MUN (mg/dl)	pH
Calculated Tank	3.87	3.62	4.65	107	15	

**Figure 1:** Somatic Cell Count (SCC) Trend

The average Somatic Cell Count (SCC) and the trend over time is shown in the graph. The ideal is a herd with an average count of less than 200 000. A higher SCC results in lower milk production and possible udder infection. If animals with high SCC can be identified early, they can be treated in time, thereby preventing production loss and permanent udder damage resulting in lower herd life.




Figure 2: Percentage (%) of Cows with Somatic Cell Counts (SCC) more than 200k and 750k, respectively, per Test-day

The graph presents the trend of the % of cows with Somatic Cell Counts (SCC) more then 200k and 500k, respectively. The ideal is to have as few cows as possible with a SCC of more than 200K and none with a SCC of more than 750k.




Figure 3: Somatic Cell Count (SCC) Monitor Trend

The graph shows the percentage of cows within certain Somatic Cell Count (SCC) levels. Ideally, at least 64% of the cows of a herd should have a SCC of less than 100 000 and 80% a SCC of less than 250 000. The danger zone is indicated in red, where ideally no cows should be indicated.




Figure 4: Average Somatic Cell Counts (SCC, x1000/ml) per Day In Milk (DIM)

This graph indicates the average Somatic Cell Counts (SCC, x1000/ml) per day in milk (DIM) for the complete herd, as well as for Lactation 1, 2, 3 and 4+ cows, separately.




Figure 5: Somatic Cell Count (SCC) Distribution for Test-day (2024-12-11)

The graph depicts the number of animals within certain levels of Somatic Cell Counts (SCC). If the cows with SCCs higher than 600 000 can be identified early, they can be treated or even culled from the herd, preventing economic losses due to penalties paid for high SCCs in the milk.

Table 7: Somatic Cell Count (SSC) distribution and Milk Loss Monitor 2024-12-11

SSC Levels	< 100k	100k - 200k	200k - 500k	500k - 800k	800k - 1 500k	> 1 500k
Number Cows	9	1	2	0	0	0
% Cows	75.0 %	8.3 %	16.7 %	0.0 %	0.0 %	0.0 %
Milk Quality	Excellent	Very Good	Needs Improvement	Poor	Serious	BAD
% Milk Loss	< 2 %	2 - 4 %	4 - 8 %	8 - 10 %	10 - 35 %	> 35 %
% Quarters Infected	< 1 %	1 - 6 %	6 - 16 %	16 - 26 %	26 - 48 %	> 48 %

The average Somatic Cell Count (SCC) on 2024-12-11 for the herd, was 93 000. This figure places the herd in the "Dark Green" SCC Zone and the milk quality is Excellent. Milk loss due to SCC is less than 2 %.

Udder Health: Very little mastitis is present. Monitor individual cows.

Table 8: Cows Under Pressure (SCC > 500k and Lactose % < 4.6)

Cow ID	Name	SCC	Lactose %	DIM	Parity	Production
--------	------	-----	-----------	-----	--------	------------

Table 9: Cows with Somatic Cell Counts > 200k (Status: N=New -, C=Chronic -, R=Recurring - and CR=Chronic and Recurring Infections)

Cow ID	Name	SCC 2024-12-11	SCC 2024-10-18	SCC 2024-09-27	Production	DIM	Parity	Status
HRG507	AWARD LOBELLIA I	422	78	22	28	132	7	R
HRD1166	BRIAR	256	149	48	20.6	320	9	R

Table 10: Possible Ketosis Cases; Cows under 100 days in milk (Prot. Fat Ratio < 0.68 for Jerseys and < 0.74 for other Breeds)

Cow ID	Name	Production	DIM	Parity	Prot % to Fat %	Breed
--------	------	------------	-----	--------	-----------------	-------

Table 11: Possible Acidosis Cases (Protein % > Butterfat %)

Cow ID	Name	Production	DIM	Parity	Prot % to Fat %
HRD3102	VALENTINO MERIDA	18.4	147	9	1.15
HRG408	VALENTINO BOKKIE I	20	244	8	1.15
HRG002	GLISTEN BLINKERS	29.6	23	3	1.09
HRG206	ENDURANCE BONTROK	18	136	1	1.05

Table 12: Dry Cows with a Dry Period more than 55 Days

Cow ID	Name	Breed	Last ICP	Avg. ICP	No. ICPs	Days Since Last Calf	Days Open
HRD 040011	HYDEAWAY XF FIONA	JSE	473	453	8	2334	2077
HRG801	DURABLE BLINKERS I	JSE	462	464	2	864	567
HRG120	WHISTLER BLINKERS II	JSE	-	-	-	562	243
HRG119	WHISTLER BLINKERS I	JSE	-	-	-	435	114
HRG005	GLISTEN BLINKERS III	JSE	386	386	1	475	106

Table 13: Cows with an active Inter-Calving Period (ICP) > 400 days

Cow ID	Name	Breed	Last ICP	Avg. ICP	No. ICPs	Days Since Last Calf	Production
HRD 040011	HYDEAWAY XF FIONA	JSE	473	453	8	2334	-
HRG801	DURABLE BLINKERS I	JSE	462	464	2	864	-
HRG120	WHISTLER BLINKERS II	JSE	-	-	-	562	-
HRG005	GLISTEN BLINKERS III	JSE	386	386	1	475	-
HRG119	WHISTLER BLINKERS I	JSE	-	-	-	435	-