

Boerboel Genetic Evaluation

SA Stud Book
February 2026

MR C. BOONZAAIER

8 HIGH ROAD
WOODLANDS
CAPE TOWN
7785



Kennel Number: 706816



Estimated Breeding Values (EBVs)

Litter size and conformation traits are influenced by many genes, with every gene making a small contribution to the expression of the trait. These traits are, however, also influenced by physiological and physical environmental factors, for example the age and sex of the animal, the feeding and management the animal received and year x season effects. It is therefore not possible to know what the animal's genetic potential for the trait is by merely using the measurement.

With BLUP (*Best Linear Unbiased Prediction*) methodology an animal's genetic potential is predicted by making use of the genetic parameters (heritabilities and genetic correlations) of the breed, pedigree information, as well as the measurements of the animal as deviation from the average of the animals that were measured with the animal. Adjustments are made for differences in sex, age, classifier, management and year x season effects to ensure that fair comparisons are made amongst the animals. These predictions are called **EBVs (Estimated Breeding Values)** which indicates the animal's genetic potential for the specific trait and predicts the type of progeny that the animal will breed.

Pedigree information determines the genetic level at which the animal enters the genetic evaluation, it links the information of all relatives together and it is used to determine the genetic ranking of the breed. For young animals that were not yet measured, mid-parent EBVs are estimated, which are the average of the EBVs of the sire and dam. These breeding values are still inaccurate and can change a lot when more information (measurements) is included in subsequent genetic evaluations.

Breeding values therefore enables a breeder to compare his/her animals genetically to one another, as well as to all other animals of the breed. This Report gives a summary of the genetic levels of the active males and females of the relevant breeder in comparison to that of the breed for linearly classified traits and litter size. The genetic change that occurred over time can also easily be seen.

Breeding Value Indices (EBVi)

For easy interpretation, EBVs are standardised and expressed relative to the average (100) of the active animals of the breed, using 12 as standard deviation (SD). This means that an EBV of 100 is equal to the active average of the breed, and more or less 50% of the animals will have EBVi's of between 90 and 110, which indicates an average animal. Only 0.5% of the EBVi's will be lower than 64 and 0.5% will be higher than 136. Higher EBVi's indicate better genetic potential for the trait.

Accuracy

EBVs are published with an **accuracy value (%)**, which indicates how much info was available for the estimation of the EBV and with how much trust it can therefore be applied. Factors that influence the accuracy include heritability, completeness and correctness of pedigree information, number of own and measured relatives and the size of the group in which the animal was measured. When an accuracy of 85% and more is reached, the EBV should not change significantly with the inclusion of more information in subsequent genetic evaluations.

Kennel and Breed Averages: The different traits' averages of the breeder's active animals are compared to those of the breed in the Kennel and Breed Average Graphs and Tables. The breeder can therefore easily see for which traits the animals are above or below the average of the breed.


Genetic Trends: The Genetic Trends indicate the genetic change that occurred over time in the animals bred by the breeder compared to that of the breed. These are calculated as the average of the breeding values of all measured animals bred by the breeder, per year of birth.

Inbreeding: Inbreeding is the mating of related animals and is widely viewed as being harmful. The danger of inbreeding lies there in that it can gradually decrease fertility and survivability – a phenomenon known as inbreeding depression. It can cause lower conception rates, more abortions, more stillborn and weak offspring and a higher susceptibility to diseases. Indications in dogs include a low litter size as well as problems relative to fertility, such as invisible heat, behavioural problems when mating and low sperm quality. Overall, reduced litter size most likely reflects inbreeding depression in several traits, both physiological and behavioural, as well as a high frequency of non-viable embryos due to lethal alleles. Generally, the effects of inbreeding can become noticeable at a value of 6.25%, therefore it is generally recommended that individuals should have inbreeding coefficients of less than 6.25%. Breeders should therefore avoid matings causing inbreeding of more than 6.25% in the progeny. The theoretical maximum inbreeding coefficient is 50%. However, it should be noted that not all inbred animals show signs of inbreeding depression.

The data used in this evaluation was obtained from LOGIX from linear classifications that were done by the Breeder Society. All possible attempts have been made to present the information correctly. SA Stud Book does not accept responsibility for the use and interpretation of the information presented on animals in this publication.

Genetic Levels

Kennel Average relative to Breed Average, where Breed Average is indicated as 0



Average Breeding Value Indices (EBVi) where Breed Average = 100

Trait	All Animals	Males	Females
Litter size (LS)	107	104	109
Movement (Mv)	100	100	100
Temperament (Tmp)	105	116	100
Coat (Ct)	102	111	97
General Appearance	111	129	102
Head & Face (H&F)	112	127	105
Front Quarter (FQ)	111	120	107
Centre (C)	105	112	102
Hind Quarter (HQ)	101	108	98
Final Score (FS)	108	121	102
Hip Health (HH)	111	118	107
Elbow Health (ELM)	101	100	102

Inbreeding

Notes

Inbreeding primarily leads to decreased fertility, low survivability of pups and low disease resistance.

Inbreeding coefficients are calculated for all animals of the breed and indicates the probability that two genes of a pair are the same due to a common ancestor in the pedigree of the animal.

The adverse effects of inbreeding can be observed from 6.25% onwards.


It is therefore highly recommended to record pedigree information as accurately. Inbreeding is dependent on pedigree information. Conventional methods for the calculation of inbreeding coefficients usually assume that unknown parents are unrelated, which is usually far from the truth. Inbreeding coefficients are therefore conservative indicators of the level of inbreeding.

Incomplete or incorrect pedigree information can be reported to SA Stud Book via the Breed Society.

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Rate of Inbreeding




Inbreeding Levels


	All Animals		Males		Females		Sires		Dams	
	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed
Number	3	49007	1	25141	2	23857	3	3371	3	6287
Average Inbreeding	9.47	8.49	7.14	8.46	10.64	8.53	5.29	8.01	8.31	8.00
Number inbred	3	48727	1	24990	2	23728	3	24990	3	23728
Number >6.25% Inbr.	3	30600	1	15537	2	15058		15537	2	15058

Genetic Trends


Litter Size




Movement




Temperament




Coat



General Appearance




Head & Face




Genetic Trends


Front Quarter



Centre




Hind Quarter




Final Score & Health Traits


Final Score



Hip Health



Elbow Health



Estimated Breeding Value Indices (EBVi's)

Males

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :	Appraisal	Health		Progeny with litters
Name	Sire	Dam	%	Hips	Elbows	Litter Size	Move-	Tempe-	Coat	General	Head &	Front	Centre	Hind	Final	Progeny	Progeny	Hips	Elbows	
	BYr	Dam		Acc.	Acc.	Acc.	ment	rament		Appear.	Face	Quarter		Quarter	Score	Litters	Kennels			
PIONEER MIDAS		MIDDELPOS BBD BRUCE ALMIGHTY		7.14	118	100	104	100	116	111	129	127	120	112	108	121				
	2024	MALAKAS HONEY			5	5	7	42	15	53	41	53	42	44	46	50				

Estimated Breeding Value Indices (EBVi's)

Females

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information

Inbreed

Health

Fertility

General

Appearance

Progeny :

Appraisal

Health

Progeny with litters

Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	Progeny with litters
GROOT HART GYPSY		AFRIKA TOMMELOM	14.86	117	107	123	95	105	102	112	115	116	109	105	111	8	2	1			1
	2019	AFRIKA MIA 9		5	13	28	48	24	61	47	60	49	50	52	57						
PIONEER MEDUSA		LOWEN KAROOLAND SAM	6.42	98	96	95	106	96	92	92	95	97	94	92	93	6	1	1			
	2020	WALSH PEPPER		5	5	5	43	9	60	42	59	43	46	49	54						

Estimated Breeding Value Indices (EBVi's)

Sires

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Sire Information		Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels	Hips	Elbows	
LOWEN KAROOLAND SAM		MIDDELPOS BBD JAPIE	4.66	90	108	93	89	83	101	96	90	96	95	81	87	12	3	5		4
	2013	AFRIKA KARLA II		5	5	20	50	22	67	49	66	51	53	56	62		4			2
AFRIKA TOMMELOM		MIDDELPOS BBD ALPHA	5.38	128	111	128	92	108	72	104	114	122	110	107	108	80	22	32	4	19
	2015	AFRIKA BLITZY		43	54	61	75	55	85	74	85	75	77	79	82		26		4	3
MIDDELPOS BBD BRUCE ALMIGHTY		AFRIKA LEO 5	5.84	127	106	106	86	104	108	119	125	112	109	111	117	48	8	27	2	16
	2017	MIGHTY AFRIKA ZELLIE		32	49	37	69	44	82	68	81	70	72	74	78		18		2	5

Estimated Breeding Value Indices (EBVi's)

Dams

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe- r ament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
WALSH PEPPER		ATKINS GRIDLOCK	7.18	105	85	96	116	104	104	90	100	96	91	100	99	15	2	4			3
	2018	WALSH WINNIE 3		5	5	13	50	23	68	50	67	51	54	56	62			4			1
AFRIKA MIA 9		AFRIKA RAMBOMAX	11.97	106	103	118	103	106	108	118	116	110	112	108	114	8	1	4			3
	2018	AFRIKA KONTESSA		23	38	44	53	33	65	52	64	53	55	57	61			3	1	1	1
MALAKAS HONEY		ELEVATION GUNNER	5.80	109	93	103	115	128	114	139	129	128	115	104	125	13	2	9			2
	2021	DELATRO CARA		5	5	17	52	22	69	51	68	53	55	58	64			6			1

