

Boerboel Genetic Evaluation

SA Stud Book
February 2026

MR J. TAYLOR

6 FROSTY ACRES

SHAWNEE

OK 24801

USA



Kennel Number: 706560



Estimated Breeding Values (EBVs)

Litter size and conformation traits are influenced by many genes, with every gene making a small contribution to the expression of the trait. These traits are, however, also influenced by physiological and physical environmental factors, for example the age and sex of the animal, the feeding and management the animal received and year x season effects. It is therefore not possible to know what the animal's genetic potential for the trait is by merely using the measurement.

With BLUP (*Best Linear Unbiased Prediction*) methodology an animal's genetic potential is predicted by making use of the genetic parameters (heritabilities and genetic correlations) of the breed, pedigree information, as well as the measurements of the animal as deviation from the average of the animals that were measured with the animal. Adjustments are made for differences in sex, age, classifier, management and year x season effects to ensure that fair comparisons are made amongst the animals. These predictions are called **EBVs (Estimated Breeding Values)** which indicates the animal's genetic potential for the specific trait and predicts the type of progeny that the animal will breed.

Pedigree information determines the genetic level at which the animal enters the genetic evaluation, it links the information of all relatives together and it is used to determine the genetic ranking of the breed. For young animals that were not yet measured, mid-parent EBVs are estimated, which are the average of the EBVs of the sire and dam. These breeding values are still inaccurate and can change a lot when more information (measurements) is included in subsequent genetic evaluations.

Breeding values therefore enables a breeder to compare his/her animals genetically to one another, as well as to all other animals of the breed. This Report gives a summary of the genetic levels of the active males and females of the relevant breeder in comparison to that of the breed for linearly classified traits and litter size. The genetic change that occurred over time can also easily be seen.

Breeding Value Indices (EBVi)

For easy interpretation, EBVs are standardised and expressed relative to the average (100) of the active animals of the breed, using 12 as standard deviation (SD). This means that an EBV of 100 is equal to the active average of the breed, and more or less 50% of the animals will have EBVi's of between 90 and 110, which indicates an average animal. Only 0.5% of the EBVi's will be lower than 64 and 0.5% will be higher than 136. Higher EBVi's indicate better genetic potential for the trait.

Accuracy

EBVs are published with an **accuracy value (%)**, which indicates how much info was available for the estimation of the EBV and with how much trust it can therefore be applied. Factors that influence the accuracy include heritability, completeness and correctness of pedigree information, number of own and measured relatives and the size of the group in which the animal was measured. When an accuracy of 85% and more is reached, the EBV should not change significantly with the inclusion of more information in subsequent genetic evaluations.

Kennel and Breed Averages: The different traits' averages of the breeder's active animals are compared to those of the breed in the Kennel and Breed Average Graphs and Tables. The breeder can therefore easily see for which traits the animals are above or below the average of the breed.


Genetic Trends: The Genetic Trends indicate the genetic change that occurred over time in the animals bred by the breeder compared to that of the breed. These are calculated as the average of the breeding values of all measured animals bred by the breeder, per year of birth.

Inbreeding: Inbreeding is the mating of related animals and is widely viewed as being harmful. The danger of inbreeding lies there in that it can gradually decrease fertility and survivability – a phenomenon known as inbreeding depression. It can cause lower conception rates, more abortions, more stillborn and weak offspring and a higher susceptibility to diseases. Indications in dogs include a low litter size as well as problems relative to fertility, such as invisible heat, behavioural problems when mating and low sperm quality. Overall, reduced litter size most likely reflects inbreeding depression in several traits, both physiological and behavioural, as well as a high frequency of non-viable embryos due to lethal alleles. Generally, the effects of inbreeding can become noticeable at a value of 6.25%, therefore it is generally recommended that individuals should have inbreeding coefficients of less than 6.25%. Breeders should therefore avoid matings causing inbreeding of more than 6.25% in the progeny. The theoretical maximum inbreeding coefficient is 50%. However, it should be noted that not all inbred animals show signs of inbreeding depression.

The data used in this evaluation was obtained from LOGIX from linear classifications that were done by the Breeder Society. All possible attempts have been made to present the information correctly. SA Stud Book does not accept responsibility for the use and interpretation of the information presented on animals in this publication.

Genetic Levels

Kennel Average relative to Breed Average, where Breed Average is indicated as 0



Average Breeding Value Indices (EBVi) where Breed Average = 100

Trait	All Animals	Males	Females
Litter size (LS)	109	103	111
Movement (Mv)	110	107	112
Temperament (Tmp)	108	104	110
Coat (Ct)	104	101	106
General Appearance	104	106	103
Head & Face (H&F)	111	110	112
Front Quarter (FQ)	115	112	116
Centre (C)	113	107	116
Hind Quarter (HQ)	111	102	115
Final Score (FS)	113	109	116
Hip Health (HH)	106	106	107
Elbow Health (ELM)	98	101	97

Inbreeding

Notes

Inbreeding primarily leads to decreased fertility, low survivability of pups and low disease resistance.

Inbreeding coefficients are calculated for all animals of the breed and indicates the probability that two genes of a pair are the same due to a common ancestor in the pedigree of the animal.

The adverse effects of inbreeding can be observed from 6.25% onwards.


It is therefore highly recommended to record pedigree information as accurately. Inbreeding is dependent on pedigree information. Conventional methods for the calculation of inbreeding coefficients usually assume that unknown parents are unrelated, which is usually far from the truth. Inbreeding coefficients are therefore conservative indicators of the level of inbreeding.

Incomplete or incorrect pedigree information can be reported to SA Stud Book via the Breed Society.

Inbreeding Levels for Specific Matings

Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Rate of Inbreeding




Inbreeding Levels


	All Animals		Males		Females		Sires		Dams	
	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed	Kennel	Breed
Number	6	49007	2	25141	4	23857	3	3371	2	6287
Average Inbreeding	7.74	8.49	7.04	8.46	8.10	8.53	10.60	8.01	8.82	8.00
Number inbred	6	48727	2	24990	4	23728	3	24990	2	23728
Number >6.25% Inbr.	5	30600	1	15537	4	15058	3	15537	2	15058

Genetic Trends


Litter Size




Movement




Temperament




Coat



General Appearance




Head & Face




Genetic Trends


Front Quarter



Centre



Hind Quarter




Final Score & Health Traits

Final Score



Hip Health



Elbow Health



Estimated Breeding Value Indices (EBVi's)

Males

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe- rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny	Litters	Progeny Kennels	Hips	Elbows	
BARBARIAN JAGGER I		BARBARIAN MAROTTI	6.13	108	101	102	108	99	97	108	105	108	102	96	102	34	6				
	2022	AFRIKA JADE		5	5	6	40	5	57	39	56	41	44	46	52						
TX TAYLOR DRISCO		BRANDVLEI MARINO	7.94	105	101	104	106	110	106	105	115	116	113	109	115						
	2024	BARBARIAN NALA		5	5	12	27	5	41	26	40	27	30	32	36						

Estimated Breeding Value Indices (EBVi's)

Females

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :	Appraisal	Health		Progeny with litters
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move-ment	Tempe-rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels	Hips	Elbows	
TX TAYLOR ROGUE	2023	PETERBUILT ZEUS III	8.25	108	93	119	117	109	105	104	112	116	116	116	116					
		BARBARIAN NALA		5	5	7	30	5	49	29	48	31	34	37	43					
TX TAYLOR RAMSAY	2023	PETERBUILT ZEUS III	8.25	108	93	119	118	110	105	99	105	117	122	127	118					
		BARBARIAN NALA		5	5	7	30	5	49	21	48	31	34	37	43					
TX TAYLOR DAISY	2024	BRANDVLEI MARINO	7.94	105	101	104	106	110	106	105	115	116	113	109	115					
		BARBARIAN NALA		5	5	12	27	5	41	26	40	27	30	32	36					
TX TAYLOR YRSA	2024	BRANDVLEI MARINO	7.94	105	101	104	106	110	106	105	115	116	113	109	115					
		BARBARIAN NALA		5	5	12	27	5	41	26	40	27	30	32	36					

Estimated Breeding Value Indices (EBVi's)

Sires

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Sire Information			Inbreed	Health		Fertility	General			Appearance						Progeny :	Appraisal	Health	Progeny
Name	BYr	Sire Dam	%	Hips Acc.	Elbows Acc.	Litter Size Acc.	Move- ment	Tempe- rament	Coat	General Appear.	Head & Face	Front Quarter	Centre	Hind Quarter	Final Score	Progeny Litters	Progeny Kennels	Hips Elbows	with litters
BARBARIAN MAROTTI		BARBARIAN KONAN	12.51	98	94	98	111	103	104	115	121	111	108	95	111	17	2	6	3
	2016	BRAVAKKER BEATRICE		5	5	6	48	7	66	47	65	49	51	54	60		6		1
BRANDVLEI MARINO		MES BENOIT	10.93	103	100	79	102	106	116	108	118	108	110	109	115	30	7	1	
	2020	MES GULIETTA		5	5	27	53	28	68	52	67	54	56	58	63		1		
PETERBUILT ZEUS III		AFRIKA DROGO	8.36	110	84	108	119	109	108	105	111	116	119	123	119	20	2	3	
	2021	AFRIKA ZIMBIE		5	5	22	46	18	62	44	61	46	49	51	56		2		

Estimated Breeding Value Indices (EBVi's)

Dams

Inbreeding >6.25% is high

< 80

>= 100

>= 120

Animal Information			Inbreed	Health		Fertility	General			Appearance						Progeny :		Appraisal	Health		Progeny with litters
Name		Sire	%	Hips	Elbows	Litter Size	Move-	Tempe-r	Coat	General	Head &	Front	Centre	Hind	Final	Progeny	Litters	Progeny Kennels	Hips	Elbows	
	BYr	Dam		Acc.	Acc.	Acc.	ment	ament		Appear.	Face	Quarter		Quarter	Score						
AFRIKA JADE		AFRIKA SAMBA	10.92	117	107	106	105	83	106	88	85	95	93	107	93	17	2	6			3
	2020	MES MISHA II		5	14	36	54	30	69	54	68	55	57	60	64						
BARBARIAN NALA		BLACK IRON GOTTI	6.73	107	102	129	110	114	96	102	112	125	116	108	114	35	3	3			
	2021	BARBARIAN BERETTA		5	5	39	38	5	55	37	54	39	41	44	49						

