

KALAHARI RED GENETIC REPORT

04 September 2026



Kalahari Red Club

WATERBERG KALAHARI REDS

FDM : 303



**CERTIFICATE
OF QUALITY**

Valid up to
October 2029

General Information

This **Genetic Report** of your flock enables you to determine the genetic merit and potential of the animals in your herd relative to that of the breed. An animal's performance is determined by two factors, namely the environment and genetics. Good nutrition and management can improve performance, but an animal cannot perform better than its genetic potential. Superior performance due to the environment cannot be carried over to the next generation. It is therefore extremely important not to neglect genetic progress of your flock. Due to the interaction between environment and genetics, it is difficult to identify genetically superior animals on performance data. Estimated Breeding Values (EBVs) are therefore predicted for every animal to determine its genetic potential, as well as its worth as a parent for the next generation. Through the use of EBVs, animals can be compared fairly **within the breed**, therefore across herds, years and production systems. Results presented in this report allow the breeder to identify the underlying genetics of the animals.

Genetic parameters

Genetic parameters (heritability and genetic correlations) are estimated on a breed-basis and are required for the prediction of breeding values. The **heritability** refers to the portion of the variation of a trait in a population that is due to transferrable genetics (the rest of the variation can be attributed to the environment and non-transferrable genetics). Some traits are genetically correlated, which means that a change in one trait will lead to changes in other traits. The strength of this association amongst traits is referred to as a **genetic correlations** and can be desirable or undesirable.

Breeding Values

The prediction of EBVs makes use of performance recordings of the animal as well as those of its relatives, pedigree information and genetic parameters. The deviation of the animal's measurement relative to the average of its contemporary group, forms the basis for identifying the animal's genetic potential for the trait. A **contemporary group** is a group of animals that have been exposed to the same environmental and biological effects, including herd, year, sex and age effects. After adjusting for differences in these effects, animals can fairly be compared and the difference in performance that still persist are then assumed to be due to differences in the genetic merit of the animals.

The EBV predicted for an animal at birth and until it has been measured, is the average EBV of its parents, since the animal received half of its genes from each parent. This mid-parent value makes the assumption that the same half of the parents' genes are carried over to all their offspring. In reality, each offspring receives a random sample of each parent's genes,

therefore mid-parent EBVs are still of low accuracy, but it does indicate the genetic level that can be expected due its parental history. An animal's true genetic merit can be predicted more accurately as more information becomes available, eg. when the animal has been measured or when the number of relatives with measurements increases. Estimated breeding values therefore change over time as more information becomes available. EBVs of animals are published with an **accuracy** (%), which indicates how much information was available for the estimation of breeding values and therefore the confidence with which a breeder can use it. Factors that influence the accuracy of an EBV are the heritability of the trait, the number of measured relatives, the size of the contemporary group, the precision with which the measurements were taken and the completeness of pedigree information. An EBV with an accuracy of 85% or more is less likely to change significantly with subsequent genetic analyses.

Interpretation of breeding values

EBVs are expressed around a **base year** (5 years ago), which means that the average breeding value of the base year is set as zero and all breeding values are expressed relative to this average. An animal with an EBV of +1kg for weaning weight is therefore genetically superior by 1kg compared to an average animal born in the base year. EBVs therefore express genetic merit of animals relative to each other. The true performance of an animal cannot be calculated from the EBV, but it enables the breeder to know which animals will make a better contribution to the next generation. For easier interpretation EBVs are also expressed as breeding value indices. For these indices the average value of the **active population** is set as 100 with a standard deviation of 12. An index of 76 therefore indicates that the animal is 2 standard deviations below the average and an index of 112 indicates that the animal is 1 standard deviation above the average of the active population for the specific trait. Please note that negative EBVs are favourable for Inter-Lambing Period and Age at First Lambing, therefore shorter inter-lambing periods and younger ages at first lambing indicates better fertility. However, breeding value indices are expressed in the same way for all traits, therefore higher indices indicates desirability for the traits and lower indices are undesirable. This makes it easy to identify superior animals for all traits.

The data used in this evaluation has been obtained from LOGIX, as supplied by breeders. Everything possible had been done to present the information correctly. SA Stud Book does not take any responsibility for the use and interpretation of the information from this publication.

Selection Indices

It is easy to select on a single trait, but that is not the best approach to follow. Selecting on only one trait can lead to the deterioration of performance in other traits due to genetic correlations. There are different traits that are important, some being more important than others. A selection index combines multiple traits, each with an economic weight to take the different levels of importance into account. EBVs for different traits cannot simply be added together, since the unit of measurement and range of the values differ and in some cases negative values are favoured. These factors are accounted for in the compilation of a selection index. An overall selection index includes EBVs of all the traits of economic importance, while other selection indices are more focused on specific breeding objectives. Selection indices are expressed around 100, with 100 being the average of the active population, with a standard deviation of 12. A selection index of 124 is therefore 2 standard deviations above the average and an index of 88 is one standard deviation below the average of the active animals of the population.

Two Indices are constructed for the Kalahari Red Goat breed, namely the Logix Merit Index and the Reproduction Index. These indices, their purpose and the contribution of the different traits to the indices are summarised as follows :


Logix Merit Index (LMI)

This index combines all the traits to select for a balanced, profitable animal that is fertile and delivers a desirable level of production while not reaching very large adult weights. Very large animals require more feed to meet their maintenance requirements, which makes them more expensive to keep and more likely to struggle during difficult times. Since large animals in a breeding flock are unfavourable, a penalty is placed on Post-Wean Weight EBV (indicated by the negative value) in the index. Selection based on the LMI will lead to breeding of balanced animals that will have the genetic potential to wean at a heavier weight (both due to its own potential and having dams that produce sufficient milk), lambs at an early age, lambs regularly, give multiple lambs and has lower maintenance requirements. This value is especially important for sustainability, which requires the selection of good, productive replacement animals for the flock.

The LMI is constructed as follows :

$$LMI = (0.0675 * AFL) + (0.1125 * ILP) + (0.264 * Wean Dir) + (0.176 * Wean Mat) + (0.27 * N_Lambs_Weaned) + (0.11 * Post-Wean Dir)$$

Schematically the LMI can be explained as follows :




Reproduction Index (RI)

This index includes Weaning Maternal), Number of Lambs Weaned, Age at First Lambing and Inter-lambing. This is especially an important index to consider in the development of a dam line.

The RI is constructed as follows :

$$(0.2 * AFL) + (0.2 * ILP) + (0.3 * Wean Mat) + (0.3 * N_Lambs_Weaned)$$

Schematically it can be explained as follows :



It is always important to consider the traits that contribute to the index individually when selection of animals is based on indices. It sometimes happen that an animal receives a good index due to its exceptional genetic merit of a specific trait, while some of the other traits of the animal in the index might be at undesirable levels for the specific flock's breeding objective, but masked in the index due to the exceptionality of that trait. A sound procedure is to rank animals based on the relevant selection index and then eliminate animals from the list that do not conform towards the flock's breeding objectives when considering the individual traits. Make use of all the available information.

Genetic Levels

Breeding Values

Eienskap	Flock**			Breed		
	All	Rams	Ewes	All	Rams	Ewes
Weaning Weight Direct	0.18	0.34	0.12	0.10	0.13	0.09
Weaning Weight Maternal	-0.06	-0.15	-0.03	-0.02	-0.02	-0.03
Weaning Weight Combined	0.03	0.03	0.03	0.03	0.04	0.02
Post-wean Weight	3.97	4.66	3.70	3.97	4.04	3.94
Number Lambs Weaned	0.28	0.18	0.32	0.35	0.35	0.35
Total Weight Weaned	0.29	0.31	0.28	0.29	0.31	0.27
Age at First Lambing	-1.47	0.52	-2.23	-0.26	-0.27	-0.25
Inter-lambing Period	-0.49	-2.20	0.17	0.28	0.14	0.35
Selection Indices						
Logix Merit Index	99	99	98	98	98	98
Reproduction Index	98	97	98	97	97	97
Inbreeding						
Inbreeding (%)	3.73	3.18	3.94	5.70	5.98	5.58
Number of Animals						
Number of Animals	18	5	13	12707	3920	8764

** Averages of all active animals in the flock.

Breeding Value Indices

Trait	Flock **		
	All	Rams	Ewes
Weaning Weight Direct	100.1	102.3	99.3
Weaning Weight Maternal	96.4	94.5	97.1
Weaning Weight Comb	96.9	96.8	96.9
Post-wean Weight	98.8	103.4	97.0
Number Lambs Weaned	98.7	98.1	98.9
Total Weight Weaned	97.8	97.9	97.8
Age at First Lambing	103.7	100.9	104.7
Interlambing Period	98.6	102.4	97.2
Selection Indices			
Logix Merit Index	98.6	99.4	98.3
Reproduction Index	97.9	96.8	98.4
Inbreeding			
Inbreeding (%)	3.73	3.18	3.94
Number of Animals			
Number of Animals	18	5	13

** Averages of all active animals in the flock.

Inbreeding

NOTES

- Inbreeding primarily leads to decreased fertility and low survivability of lambs.
- The effects of inbreeding can be observed from 6.25% onwards.
- It is therefore highly recommended to record pedigree information as accurately and completely as possible and to use a selection programme for selection of unrelated rams.
- Incomplete or incorrect pedigree information can be reported to SA Stud Book. Inbreeding is dependent on pedigree information. Conventional methods for the Calculation of inbreeding coefficients usually assume that unknown parents are unrelated. However, this is usually far from the truth. An algorithm is used for the calculation of inbreeding, where the average inbreeding coefficient is assigned to unknown parents based on the generation in which they were born. This result is a more accurate reflection of inbreeding of the herd and breed.

Inbreeding Levels for Specific Matings


Progeny with Parents :	25%
Full Brother x Full Sister :	25%
Half Brother x Half Sister :	12.5%
One Common Grandparent :	6.25%
One Common Great-Grandparent :	3.125%

Level and Rate of Inbreeding

Year of Birth	Level* (%)	Rate (% / year)
2023 - 2026	2.93	0.12
Recommended:	< 6.25	< 0.1 - 0.2

*Of the last year of birth

Inbreeding




Pedigree Information

	Number	%
Known Sires	14	78
Known Dams	18	100

Note: Inbreeding can be under-estimated if pedigree information is incomplete or incorrect

Genetic Windows




WWd: Weaning Weight Direct
WWm: Weaning Weight Maternal
WWc: Weaning Weight Combined
PoW: Post-Wean Weight
L-wean: Number of Lambs Weaned
TWW: Total Weight Weaned
AFL: Age at First Lambing
ILP: Inter-lambing Period
RI: Reproduction Index
LMI: Logix Merit Index

The genetic windows allow a quick overview of the genetic status of the flock compared to the breed. The values presented here are the average EBV indices of all the active animals of the flock, of the dams and sires of the active animals of the flock, as well as of the active ewes and rams younger than 2 years old. An index of 100 is equal to the average of the active measured population. Due to the higher selection intensity on rams (selection is more strict since a much smaller percentage of young animals are retained in the flock), the expectation is for the average index of the sires to be higher than that of the dams.
 Remember that the EBV index for AFL and ILP are presented in such a way that an index above 100 indicates animals that have a lower EBV for these traits.

Genetic Windows


Dams

Number: 9




Sires

Number: 6




Young Ewes

Number: 13



Young Rams

Number: 5



Genetic Trends

Genetic trends are based on measured animals that were bred by the specific breeder.

Genetic Trends

Genetic trends are based on measured animals that were bred by the specific breeder.

For AFL and ILP flock genetic trends below breed averages are desirable.

Genetic Trends of Breed

Birth year	Weaning Weight				Post-wean	Lambs Weaned		TWW		AFL		ILP		RI		LMI				
	EBV					EBV		EBV		EBV		EBV		EBV		EBV				
	Number	Dir	Mat	Comb		Number	EBV	Number	EBV	Number	EBV	Number	EBV	Number	EBV	Number	EBV			
2013	184	0.80	-0.35	0.06	68	3.89	103	0.81	184	106	186	3.64	94	-2.39	184	96	184	106		
2014	75	-0.13	-0.10	-0.17			70	0.70	75	98	115	0.65	45	-0.59	75	97	75	98		
2015	144	-0.12	-0.14	-0.20			102	3.43	98	0.48	144	98	138	-0.07	74	-1.25	144	98	144	98
2016	108	-0.26	-0.02	-0.15			87	0.34	108	97	139	3.86	94	-4.48	108	99	108	97		
2017	271	-0.02	0.16	0.15			104	0.44	271	98	176	1.01	127	-0.49	271	99	271	96		
2018	168	-0.35	0.24	0.06			202	0.31	168	92	268	3.02	208	-2.21	168	101	168	92		
2019	293	-0.23	0.14	0.03			257	0.35	293	97	358	0.34	271	0.42	293	100	293	96		
2020	637	0.18	0.05	0.14	14	0.00	284	0.62	637	101	634	-0.01	461	0.76	637	100	637	100		
2021	562	0.00	0.00	0.00			274	-0.01	562	96	749	0.00	513	0.00	562	97	562	95		
2022	1120	-0.22	0.11	0.00			377	3.43	302	0.16	1114	97	896	0.23	522	-0.21	1114	100	1114	97
2023	1342	0.11	0.00	0.05			209	3.99	227	0.25	1342	99	805	0.42	270	0.60	1342	98	1342	98
2024	1472	0.20	0.07	0.17			164	4.47	95	-0.11	1472	101	384	-1.08	53	-2.38	1472	100	1472	101
2025	1116	0.20	0.18	0.28			394	4.65			1116	101	22	1.42			1116	102	1116	101

Genetic Trends of Flock

Birth year	Weaning Weight EBV				Post-Wean	Lambs Weaned		TWW		AFL		ILP		RI		LMI	
	Number	Dir	Mat	Comb		Number	EBV	Number	EBV	Number	EBV	Number	EBV	Number	EBV	Number	EBV

Breeding Value & Selection Index Percentiles

The percentiles indicate the ranking of the animals' EBVs and Selection Indices within the active population.

For example, if an animal's EBV for a specific trait falls between the maximum and 99% percentile of the trait, it means that this animal ranks in the top 1% of the active population, or this animal's genetic merit for the trait is better than 99% of that of the active population.

Should the EBV for a specific trait falls between the 5% and 10% percentile, it means that the animal ranks in the bottom 10% of the active population for the specific trait, or the animal's genetic merit is worse than 90% of the active population.

Note that the percentiles for Age at First Lambing and Inter-lambing Period EBVs work on the reverse scale.

All Active Animals

Trait	AVG	SA*	Min	Percentiles(%)															Max
				1	5	10	20	25	30	40	50	60	70	75	80	90	95	99	
Weaning Weight Direct	0.10	0.58	-2.61	-1.35	-0.77	-0.54	-0.31	-0.23	-0.16	-0.03	0.05	0.17	0.27	0.39	0.51	0.83	1.19	1.73	3.06
Weaning Weight Maternal	-0.02	0.41	-2.33	-0.90	-0.57	-0.45	-0.34	-0.30	-0.26	-0.18	-0.09	0.02	0.13	0.19	0.26	0.50	0.74	1.20	2.15
Weaning Weight Combined	0.03	0.48	-2.85	-1.12	-0.71	-0.54	-0.34	-0.28	-0.22	-0.08	0.03	0.13	0.23	0.29	0.36	0.60	0.87	1.30	2.36
Post-Wean Weight	3.97	1.19	-1.89	0.94	2.16	2.59	3.10	3.26	3.47	3.71	3.95	4.15	4.39	4.57	4.79	5.44	5.99	7.22	12.54
Number of Lambs Weaned	0.35	1.41	-7.50	-3.60	-1.90	-1.20	-0.50	-0.30	-0.10	0.20	0.40	0.70	0.80	1.00	1.10	1.60	2.50	4.70	13.50
Total Weight Weaned	0.29	1.17	-7.52	-3.17	-1.68	-1.10	-0.44	-0.25	-0.09	0.16	0.34	0.56	0.77	0.89	1.02	1.50	2.03	3.44	8.91
Age at First Lambing	-0.26	-7.68	56.53	18.86	11.54	8.88	5.46	4.32	3.20	1.34	0.07	-1.35	-3.26	-4.69	-6.11	-9.87	-13.80	-19.60	-37.78
Interlambing Period	0.28	-5.28	19.61	12.82	8.72	6.76	4.28	3.48	2.91	1.87	0.81	-0.54	-2.24	-3.15	-4.21	-6.85	-8.84	-12.54	-24.28
Reproduction Index	97	9	53	76	84	88	92	93	93	95	96	98	100	101	103	108	113	124	146
Logix Merit Index	98	9	45	73	83	87	92	93	95	96	98	100	101	102	104	108	112	122	147

*SD : Standard Deviation. An indicator of how far breeding values/selection indices are distributed around the average.

Breeding Value & Selection Index Percentiles

All Active Rams

Trait	AVG	SA*	Min	Percentiles (%)															Max
				1	5	10	20	25	30	40	50	60	70	75	80	90	95	99	
Weaning Weight Direct	0.13	0.58	-2.24	-1.26	-0.75	-0.51	-0.29	-0.21	-0.15	-0.02	0.07	0.19	0.31	0.43	0.55	0.86	1.21	1.70	2.81
Weaning Weight Maternal	-0.02	0.38	-1.44	-0.75	-0.52	-0.43	-0.34	-0.30	-0.25	-0.17	-0.08	0.04	0.14	0.19	0.25	0.47	0.69	1.12	1.88
Weaning Weight Combined	0.04	0.45	-1.43	-0.96	-0.64	-0.50	-0.33	-0.27	-0.21	-0.07	0.04	0.15	0.24	0.30	0.37	0.59	0.84	1.25	2.24
Post-Wean Weight	4.04	1.21	-1.77	1.42	2.31	2.63	3.12	3.28	3.47	3.73	4.00	4.22	4.48	4.67	4.91	5.49	6.15	7.30	12.54
Number of Lambs Weaned	0.35	1.31	-4.60	-3.20	-1.80	-1.10	-0.50	-0.30	-0.10	0.20	0.40	0.70	0.80	1.00	1.10	1.50	2.30	4.70	8.80
Total Weight Weaned	0.31	1.10	-4.54	-2.91	-1.53	-0.96	-0.39	-0.22	-0.08	0.15	0.34	0.59	0.80	0.92	1.02	1.45	1.95	3.29	5.84
Age at First Lambing	-0.27	-7.03	27.46	14.80	10.78	8.41	5.20	4.19	3.07	1.39	0.18	-1.27	-3.15	-4.61	-5.88	-9.01	-12.52	-19.15	-31.92
Interlambing Period	0.14	-5.04	19.27	12.69	8.29	6.32	3.96	3.17	2.66	1.58	0.49	-0.83	-2.33	-3.18	-4.11	-6.52	-8.31	-11.92	-18.90
Reproduction Index	97	8	70	77	85	89	92	93	93	95	97	98	100	101	103	108	112	124	145
Logix Merit Index	98	8	65	75	84	88	92	94	95	97	98	100	102	103	104	107	112	123	137

All Active Ewes

Trait	AVG	SA*	Min	Percentiles (%)															Max
				1	5	10	20	25	30	40	50	60	70	75	80	90	95	99	
Weaning Weight Direct	0.09	0.58	-2.61	-1.41	-0.78	-0.55	-0.32	-0.23	-0.16	-0.03	0.04	0.15	0.25	0.36	0.48	0.81	1.17	1.73	3.06
Weaning Weight Maternal	-0.03	0.42	-2.33	-0.94	-0.60	-0.46	-0.34	-0.30	-0.26	-0.18	-0.09	0.00	0.13	0.19	0.26	0.51	0.76	1.24	2.15
Weaning Weight Combined	0.02	0.49	-2.85	-1.19	-0.73	-0.56	-0.35	-0.29	-0.23	-0.09	0.02	0.11	0.23	0.29	0.35	0.60	0.88	1.37	2.36
Post-Wean Weight	3.94	1.17	-1.89	0.80	2.10	2.57	3.09	3.26	3.47	3.70	3.92	4.11	4.36	4.51	4.73	5.41	5.89	7.15	9.58
Number of Lambs Weaned	0.35	1.46	-7.50	-3.80	-2.00	-1.20	-0.50	-0.30	-0.10	0.20	0.50	0.70	0.90	1.00	1.10	1.60	2.60	4.90	13.50
Total Weight Weaned	0.27	1.21	-7.52	-3.26	-1.74	-1.16	-0.48	-0.25	-0.10	0.17	0.34	0.54	0.75	0.88	1.02	1.53	2.04	3.50	8.91
Age at First Lambing	-0.25	-7.98	56.53	19.28	11.85	9.04	5.56	4.41	3.25	1.33	0.02	-1.39	-3.32	-4.74	-6.21	-10.44	-14.24	-19.76	-37.78
Interlambing Period	0.34	-5.40	19.61	12.85	9.02	6.90	4.47	3.57	2.96	2.01	0.96	-0.41	-2.16	-3.12	-4.28	-7.01	-9.03	-12.81	-24.28
Reproduction Index	97	9	53	75	84	88	91	93	93	95	96	98	100	101	103	109	114	125	146
Logix Merit Index	98	9	45	72	83	87	92	93	94	96	98	99	101	102	104	108	113	122	147

*SD : Standard Deviation. An indicator of how far breeding values/selection indices are distributed around the average.

Rams

Inbreeding in red is high

EBV - Estimated Breeding Value

Indices >=124
Above breed average
Indices <= 76

Acc: Accuracy of the breeding value
EBVs with an accuracy below 5% are not printed

Index: Expresses the EBV around 100 where 100 is set as the average EBV of active animals

Animal Information							Selection Indices		EBV - Weights				EBV - Fertility							
Animal Comp no.			Parents Sire Dam			% In- breeding	Logix Merit Index	Reproduction Index	Wean Weight		Post-Wean	Number Weaned	TWW		AFL		ILP		Dam	
									Dir EBV Acc Index	Mat EBV Acc Index			Comb EBV Index	EBV Acc Index	EBV Acc Index	EBV Acc Index	EBV Acc Index	EPI	Times Lambd	
Active Breed Average:							100	100	0.10	-0.02	0.03	3.97	0.35	0.29	-0.26	0.28				
232 24 0004 107598708	-	-	-	2.30	86	100	-0.83 34 86	0.36 32 106	-0.05 95	2.48 26 89	-1.50 20 88	-1.37 30 85	0.86 26 101	-5.08 15 109	88	5				
303 25 0001 114390230	3	22	0149	2.80	102	93	0.73 16 108	-0.39 13 89	-0.02 96	5.48 6 109	0.40 5 99	0.58 12 100	2.55 25 98	-2.45 14 103				2		
303 25 0002 114390248	3	22	0149	2.80	102	93	0.73 16 108	-0.39 13 89	-0.02 96	5.48 6 109	0.40 5 99	0.58 12 100	2.55 25 98	-2.45 14 103				2		
303 25 0003 114390222	3	22	0149	4.00	104	99	0.54 13 105	-0.16 12 94	0.11 99	4.93 7 105	0.80 5 102	0.87 11 102	-1.68 26 104	-0.52 14 99				2		
303 25 0004 114390255	3	22	0149	4.00	104	99	0.54 13 105	-0.16 12 94	0.11 99	4.93 7 105	0.80 5 102	0.87 11 102	-1.68 26 104	-0.52 14 99				2		
Ram Flock Average: Number of rams in the flock: 5							99	97	0.34	-0.15	0.03	4.66	0.18	0.31	0.52	-2.20				

Ewes

Inbreeding in red is high

EBV - Estimated Breeding Value

Indices >=124
Above breed average
Indices <= 76

Acc: Accuracy of the breeding value
EBVs with an accuracy below 5% are not printed

Index: Expresses the EBV around 100 where 100 is set as the average EBV of active animals

Animal Information							Selection Indices			EBV - Weights				EBV - Fertility					
Animal Comp no.			Parents			% In- breedin	Logix Merit Index	Reproduction Index	Wean Weight		Comb	Post-Wean	Number Weaned	TWW	AFL	ILP	Phenotype		
			Sire	Dam					Dir	Mat									EBV
								EBV Acc Index	EBV Acc Index	EBV Index	EBV Acc Index	EBV Acc Index	EBV Acc Index	EBV Acc Index	EBV Acc Index	EPI	Times Lambd		
			Active Breed Average:				100	100	0.10	-0.02	0.03	3.97	0.35	0.29	-0.26	0.28			
232 23 0034	-	-	-	2.57		101	100	0.03	0.05	0.06	3.25	1.20	0.80	-6.11	5.02		2		
107515645	AH	21	0140					23 98	17 99	98	20 94	6 104	17 102	20 110	11 86				
232 23 0061	-	-	-	2.57		99	92	0.56	-0.48	-0.20	4.75	0.40	0.33	-0.41	0.07		2		
107515363	147	22	0005					23 105	20 87	93	5 104	7 99	18 98	17 102	14 97				
232 23 0094	-	-	-	2.57		104	102	0.19	-0.02	0.08	3.66	1.20	0.89	-8.88	3.93		2		
107515355	140	19	0033					17 100	17 97	98	7 97	6 104	14 102	22 114	15 89				
157 24 0043	157	22	0067	2.06		94	96	-0.26	-0.08	-0.21	4.22	0.40	-0.10	4.50	-1.59		0		
107565699	157	20	0016					19 94	14 96	92	13 101	8 99	15 95	14 95	8 101				
303 25 0009	3	23	0137	3.07		100	97	0.41	-0.14	0.07	4.44	0.50	0.57	2.50	-1.69		0		
114390586	232	23	0034					24 103	14 95	98	23 102	5 100	16 100	22 98	12 101				
303 25 0010	3	23	0137	3.07		100	97	0.41	-0.14	0.07	4.44	0.50	0.57	2.50	-1.69		0		
114390602	232	23	0034					24 103	14 95	98	23 102	5 100	16 100	22 98	12 101				
232 25 0060	AH	21	0115	17.60		103	105	0.00	0.38	0.38	1.56	1.50	1.28	-14.93	11.92		0		
108708942	AH	21	0052					30 98	25 106	104	23 83	8 106	23 105	30 122	18 71				
157 25 0070	104	20	0078	3.06		105	95	1.44	-0.36	0.36	5.39	-0.20	0.95	3.41	-7.01		0		
113512719	147	21	0149					47 117	37 90	103	27 108	20 96	37 103	34 97	22 113				
303 26 0004	232	24	0004	3.18		95	101	-0.32	0.17	0.01	3.07	-0.10	-0.21	-4.01	-0.58		0		
114390263	232	23	0094					19 93	18 102	97	14 93	10 96	17 94	17 107	11 99				
303 26 0005	232	24	0004	3.18		95	101	-0.32	0.17	0.01	3.07	-0.10	-0.21	-4.01	-0.58		0		
114390610	232	23	0094					19 93	18 102	97	14 93	10 96	17 94	17 107	11 99				
303 26 0006	232	24	0004	3.18		95	101	-0.32	0.17	0.01	3.07	-0.10	-0.21	-4.01	-0.58		0		
114390628	232	23	0094					19 93	18 102	97	14 93	10 96	17 94	17 107	11 99				
303 26 0011	232	24	0004	2.55		93	96	-0.13	-0.06	-0.12	3.62	-0.50	-0.49	0.22	-2.51		0		
114372154	232	23	0061					21 96	19 97	94	13 96	11 94	18 92	15 101	11 103				
303 26 0012	232	24	0004	2.55		93	96	-0.13	-0.06	-0.12	3.62	-0.50	-0.49	0.22	-2.51		0		
114372162	232	23	0061					21 96	19 97	94	13 96	11 94	18 92	15 101	11 103				
Ewe flock average: Total ewes in flock: 13							98	98	0.12	-0.03	0.03	3.70	0.32	0.28	-2.23	0.17			

Sires Used in Flock

*Inbreeding in red is high

EBV - Estimated Breeding Value

Indices >=124
Above breed average
Indices <= 76

Acc: Accuracy of the breeding value
EBVs with an accuracy below 5% are not printed

Index: Expresses the EBV around 100 where 100 is set as the average EBV of active animals

Animal Information		Parents & Progeny				Selection Indices		EBV - Weights				EBV - Fertility					
Animal ID Computer nr	% In- breedin	Sire ID Dam ID	1st Generation		2nd Generation		Logix Merit Index	Reproducti on Index	Weaning Weight		Comb EBV Index	Post-Wean EBV Acc Index	Number Weaned EBV Acc Index	TWW EBV Acc Index	AFL EBV Acc Index	ILP EBV Acc Index	DAM EPI Times Lambe
			Progeny Number Weaned	Herds	Progeny Number Weaned	Herds			Dir EBV Acc Index	Mat EBV Acc Index							
		Active Breed Average:				100	100	0.10	-0.02	0.03	3.97	0.35	0.29	-0.26	0.28		
1	104 20 0078 86421708	11.36	104 18 0057 104 17 0045	3	1	98	99	0.33 70	-0.22 54	-0.06 95	3.59 54	0.40 35	0.34 56	10.16 55	-11.04 34	123 10	
2	AH 21 0115 96037965	17.14	AH 18 0104 AH 17 1013			100	100	-0.09 38	0.17 34	0.13 99	1.58 25	1.40 11	0.92 30	-13.40 49	13.06 26	9	
3	157 22 0067 96385448	2.64	104 20 0068 157 19 0076	1	1	92	97	-0.56 38	0.11 28	-0.17 93	4.38 25	0.00 16	-0.45 29	8.25 27	-4.16 16	5	
4	3 22 0149 95709275	16.90	3 20 0135 3 20 0130			104	95	0.90 21	-0.30 18	0.15 99	6.20 12	0.30 5	0.78 16	5.52 47	-4.97 24	1	
5	3 23 0137 98890460	9.95	3 21 0072 3 20 0082	4	1	99	93	0.79 42	-0.33 21	0.07 98	5.62 41	-0.10 7	0.40 26	11.11 39	-8.39 22	2	
6	232 24 0004 107598708	2.30	- - - 104 21 0011			86	100	-0.83 34	0.36 32	-0.05 95	2.48 26	-1.50 20	-1.37 30	0.86 26	-5.08 15	88 5	
		Sire Flock Average: Number of sires used in the flock: 6				96	97	0.09	-0.03	0.01	3.98	0.08	0.11	3.75	-3.43		

Ram Selection List: LMI

Animals sorted on LMI

Inbreeding in red is high

EBV - Estimated Breeding Value

Indices >=124
Above breed average
Indices <= 76

Acc: Accuracy of the breeding value
EBVs with an accuracy below 5% are not printed

Index: Expresses the EBV around 100 where 100 is set as the average EBV of active animals

Animal Information		Parents & Progeny				Selection Indices		EBV - Weights				EBV - Fertility				
Animal ID Computer nr	% In-breedi	Sire ID Dam ID	1st Generation		2nd Generation	Logix Merit Index	Reproducti on Index	Weaning Weight		Comb EBV Index	Post-Wean EBV Acc Index	Number Weaned EBV Acc Index	TWW EBV Acc Index	AFL EBV Acc Index	ILP EBV Acc Index	
			Progeny Number Weaned	Herds	Progeny Number Weaned			Herds	Dir EBV Acc index							Mat EBV Acc Index
		Active Breed Average:				100	100	0.10	-0.02	0.03	3.97	0.35	0.29	-0.26	0.28	
1	303 25 0003 114390222	4.00	3 22 0149 232 23 0094			104	99	0.54 13 105	-0.16 12 94	0.11 99	4.93 7 105	0.80 5 102	0.87 11 102	-1.68 26 104	-0.52 14 99	
2	303 25 0004 114390255	4.00	3 22 0149 232 23 0094			104	99	0.54 13 105	-0.16 12 94	0.11 99	4.93 7 105	0.80 5 102	0.87 11 102	-1.68 26 104	-0.52 14 99	
3	303 25 0002 114390248	2.80	3 22 0149 232 23 0061			102	93	0.73 16 108	-0.39 13 89	-0.02 96	5.48 6 109	0.40 5 99	0.58 12 100	2.55 25 98	-2.45 14 103	
4	303 25 0001 114390230	2.80	3 22 0149 232 23 0061			102	93	0.73 16 108	-0.39 13 89	-0.02 96	5.48 6 109	0.40 5 99	0.58 12 100	2.55 25 98	-2.45 14 103	
5	232 24 0004 107598708	2.30	- - - 104 21 0011			86	100	-0.83 34 86	0.36 32 106	-0.05 95	2.48 26 89	-1.50 20 88	-1.37 30 85	0.86 26 101	-5.08 15 109	

Ewe Selection List: LMI

Animals sorted on LMI

Inbreeding in red is high

EBV - Estimated Breeding Value

Indices >=124
Above breed average
Indices <= 76

Acc: Accuracy of the breeding value
EBVs with an accuracy below 5% are not printed

Index: Expresses the EBV around 100 where 100 is set as the average EBV of active animals

Animal Information		Parents		Selection Indices		EBV - Weights				EBV - Fertility				
Animal ID Computer nr	% In- breedin	Sire ID Dam ID	Logix Merit Index	Reproducti on Index	Dir EBV Acc Index	Weaning Weight Mat EBV Acc Index	Comb EBV Index	Post-Wean EBV Acc Index	Number Weaned EBV Acc Index	TWW EBV Acc Index	AFL EBV Acc Index	ILP EBV Acc Index	Phenotype	
Active Breed Average:				100	100	0.10	-0.02	0.03	3.97	0.35	0.29	-0.26	0.28	
1 303 25 0003 114390222	4.00	3 22 0149 232 23 0094	104	99	13 0.54 105	12 -0.16 94	0.11 99	4.93 7 105	0.80 5 102	0.87 11 102	-1.68 26 104	-0.52 14 99		
2 303 25 0004 114390255	4.00	3 22 0149 232 23 0094	104	99	13 0.54 105	12 -0.16 94	0.11 99	4.93 7 105	0.80 5 102	0.87 11 102	-1.68 26 104	-0.52 14 99		
3 303 25 0002 114390248	2.80	3 22 0149 232 23 0061	102	93	16 0.73 108	13 -0.39 89	-0.02 96	5.48 6 109	0.40 5 99	0.58 12 100	2.55 25 98	-2.45 14 103		
4 303 25 0001 114390230	2.80	3 22 0149 232 23 0061	102	93	16 0.73 108	13 -0.39 89	-0.02 96	5.48 6 109	0.40 5 99	0.58 12 100	2.55 25 98	-2.45 14 103		
5 232 24 0004 107598708	2.30	- - - 104 21 0011	86	100	34 -0.83 86	32 0.36 106	-0.05 95	2.48 26 89	-1.50 20 88	-1.37 30 85	0.86 26 101	-5.08 15 109		

